

**RANDOMIZED TRIALS OF SECONDARY PREVENTION
PROGRAMS IN CORONARY HEART DISEASE: A SYSTEMATIC
REVIEW**

Prepared For:

Agency for Healthcare Research and Quality
US Department of Health and Human Services
540 Gaither Road
Rockville, MD 20852
<http://www.ahrq.gov>

Contract No.

Prepared By:

Alex Clark, Lisa Hartling, Ben VanderMeer, Finlay A McAlister
for the University of Alberta Evidence-based Practice Center
Edmonton, Alberta, Canada

Funding: AC and FMcA are Population Health Investigators of the Alberta Heritage Foundation for Medical Research; FMcA also holds a Canadian Institutes of Health Research New Investigator Award and the Merck Frosst/Aventis Chair in Patient Health Management at the University of Alberta.

Correspondence to: Dr. F. McAlister (Division of General Internal Medicine, 2E3.24 WMC, University of Alberta Hospital, 8440 112 Street, Edmonton, Alberta, Canada T6G 2R7 ; e-mail: Finlay.McAlister@ualberta.ca)

Table of Contents

Abstract.....	3
Introduction.....	6
Methods.....	10
Results.....	16
Discussion.....	23
Acknowledgements.....	34
References.....	35
Box 1 (Modifiable coronary risk factors).....	50
Table 1 (Description of studies included).....	51
Table 2 (Impact on all-cause mortality and recurrent myocardial infarctions).....	57
Table 3 (Impact on other endpoints).....	59
Figure 1 (Flow of trials through the selection process).....	61
Figure 2 (All-cause mortality in trials evaluating comprehensive cardiac rehabilitation).....	62
Figure 3 (All-cause mortality in trials evaluating group cardiac rehabilitation without exercise component).....	63
Figure 4 (All-cause mortality in trials evaluating individual counseling).....	64
Figure 5 (Recurrent myocardial infarctions in trials evaluating comprehensive cardiac rehabilitation).....	65
Figure 6 (Recurrent myocardial infarctions in trials evaluating group cardiac rehabilitation without exercise component).....	66
Figure 7 (Recurrent myocardial infarctions in trials evaluating individual counseling).....	67
Figure 8 (Hospitalization rates in trials evaluating secondary prevention programs).....	68
Figure 9 (Funnel Plot for all-cause mortality data).....	69
Appendix A (Search Strategy and Results).....	70
Appendix B (List of Excluded Studies and Reasons).....	73

ABSTRACT

Objective: To determine whether secondary prevention programs for patients with established coronary artery disease (CAD) improve health outcomes.

Design: Randomized clinical trials (RCTs) of secondary prevention programs in patients with CAD were identified by searching Medline 1966-2004; the Cochrane Central Register of Controlled Trials, Issue 4, 2004; Embase 1980-2004; CINAHL 1982-2004; SIGLE 1980-2004; the Cochrane Effective Practice and Organization of Care Study Registry; bibliographies of published studies, and via contact with experts in the field and references provided by the Centres for Medicare and Medicaid Services. Studies were selected and data extracted independently by 2 investigators, and summary risk ratios were calculated using both the random and fixed effects models.

Results: A total of 41 RCTs (18 281 patients with CHD) were identified. Secondary prevention programs had positive impacts on processes of care: patients randomized to these programs were more likely to be prescribed efficacious medications and 20 out of 24 trials evaluating cholesterol profiles demonstrated improvements with

these programs compared to usual care (in 12 trials the improvements were statistically significant, with effect sizes in the small to moderate range). The summary RR was 0.89 (95% CI 0.79-1.01) for all-cause mortality, but this result differed over time with a RR of 0.97 (95% CI 0.82-1.14) for 12 month all-cause mortality in the 18 trials (9192 patients, p for heterogeneity=0.90, I^2 =0%) reporting this outcome and a RR of 0.53 (95% CI 0.31-0.92) for all-cause mortality at 24 months in the 4 trials (1367 patients, p for heterogeneity=0.44, I^2 =0%) reporting this outcome. The summary RR was 0.89 (95% CI 0.77-1.04) for recurrent myocardial infarction and 0.85 (95% CI 0.78-0.93) for hospitalization rates over a median follow-up of 12 months. There were no appreciable differences between group cardiac rehabilitation programs or secondary prevention programs focusing on individual education, counseling, and supervision in the primary outcomes we considered (mortality, hospitalizations, recurrent myocardial infarctions). Fourteen of the 26 trials evaluating quality of life or functional status reported statistically significantly better outcomes in those patients exposed to the intervention programs, although the effect sizes were generally small. Only 6 of these trials reported the costs of the

intervention- in 2 cases, the interventions were cost-saving. Another 2 trials reported significant reductions in health care resource use in patients exposed to secondary prevention programs compared to controls.

Conclusions: Secondary prevention programs improve processes of care, enhance quality of life/functional status, reduce hospitalizations, and reduce long-term mortality in patients with established CAD. Although these clinical benefits are likely to reduce health care costs, there is inadequate data to conclusively comment on the cost-effectiveness of these programs and specific components contained therein.

INTRODUCTION

Although cardiovascular death rates in North America have declined over the past two decades,[1] cardiovascular disease remains the most common cause of death (38% of all deaths in the United States in 2002), hospitalization, and physician office visits (over 80 million visits in 2002) and accounts for a large portion of total health care costs in the United States (estimated direct and indirect costs for 2005 are over \$393 billion).[2] Using data from the National Health and Nutrition Examination Survey (NHANES 1999-2002), it is estimated that over 70 million Americans have one or more types of cardiovascular disease and over 13 million have known coronary artery disease (the proportion with undiagnosed disease is likely several fold higher).[2] Indeed, it is estimated that an American suffers a coronary event every 26 seconds, with 41% dying within a year.[2] Of course, coronary artery disease is not an American phenomenon and atherosclerotic cardiovascular disease is the leading cause of death worldwide.[3] A case-control study in nearly 30,000 subjects from 52 countries confirmed that 9 known coronary risk factors (Box 1) account for over 90% of the population

attributable risks for coronary disease in both men and women, in all age subgroups, and across all regions.[4]

Control of the CAD epidemic will require a multifaceted strategy targeting the 9 modifiable risk factors delineated in the INTERHEART study and including both primary prevention maneuvers (some designed for the general population and some targeting only high risk individuals) and secondary prevention programs (targeted at those with established disease).[5] Despite the abundant evidence base for CAD prevention,[6] health outcomes studies consistently demonstrate suboptimal control of cardiovascular risk factors due to gaps in the application of this evidence to clinical practice which contribute to sub-optimal patient outcomes.[7-15] Furthermore, even when some secondary prevention therapies are prescribed, patient compliance may be poor (from 43% to 75% at one year).[16,17]

Secondary prevention programs are increasingly advocated as a means to improve management and outcomes for patients with CHD. While numerous reviews have shown that cardiac rehabilitation programs improve outcomes in MI survivors,[18-22] these conclusions are based largely on trials which tested supervised exercise programs versus no exercise. As activity levels are

inversely proportional to cardiovascular mortality and exercise training confers substantial physiologic and clinical benefits,[23] it is not surprising that trials of exercise programs found positive treatment effects.

However, few of the trials included in these reviews evaluated comprehensive secondary prevention programs employing disease management approaches. Disease management has been defined as “a combination of patient education, provider use of practice guidelines, appropriate consultation, and supplies of drugs and ancillary services”.[24] To address this gap in the literature, we performed a systematic review of randomized trials of multidisciplinary disease management programs in patients with established CAD- in the 12 trials we identified (with 9803 patients), we found that multidisciplinary disease management programs improved processes of care (namely prescription of proven efficacious secondary prevention therapies) and risk factor profiles, reduced hospitalizations by 16% (95% confidence interval [CI] 6% to 24%), but did not have an appreciable impact on rates of death (RR 0.91, 95% CI 0.79-1.04) or recurrent myocardial infarction (RR 0.94, 95% CI 0.80-1.10).[25]

As current guidelines recommend that secondary prevention programs should not be restricted to supervised exercise programs but rather address the full range of modifiable risk factors,[26,27] we conducted the current systematic review to update our earlier work and to determine whether comprehensive secondary prevention programs (in contradistinction to exercise-only or similar single modality programs) prevent coronary events and/or death in patients with CAD.

METHODS

Searching for relevant studies:

We searched the following electronic databases to identify human randomized trials published in English: Medline 1966-2004; Cochrane Central Register of Controlled Trials, Issue 4, 2004; Embase 1980-2004; CINAHL 1982-2004; SIGLE 1980-2004. In order to identify recent publications, we also searched PubMed from January 2004 to December 2004 and conducted a cited reference search for our previous systematic review[25] in Web of Science (1999 to 2004). The searches (see Appendix A for listing of search strategy strings and results) were based on the following terms: case management, comprehensive health care, disease management, health services research, home care services, clinical protocols, patient care planning, quality of health care, rehabilitation, nurse led clinics, special clinics, and myocardial ischemia. To identify any studies missed by the literature searches, we hand-searched reference lists of all identified studies, as well as the reference list of a recent related review.[22] Finally, we screened references provided by the Centres for Medicare and Medicaid Services and content experts.

Selection of studies and abstraction of data:

Two of the investigators (AC and FM) independently reviewed the titles and abstracts of all citations to identify any studies reporting the impact of secondary prevention programs on death, MI, or hospitalization rates in patients with CAD (clinically manifest as angina, MI, or coronary revascularization). The full texts of all potentially relevant articles were obtained and reviewed by both investigators using pre-standardized data abstraction forms and a *priori* defined eligibility criteria. Any discrepancies were resolved by consensus.

All outcome data were extracted by AC and FM independently, and double-checked by BV. Outcomes were assigned according to the intention-to-treat principle and we accepted the definitions for each outcome used by the investigators in the primary studies. Original investigators were contacted to clarify the published data for any trials published in the past decade for which our outcomes of interest were not reported: authors for 9 of the 21 studies contacted provided further data.

Studies were excluded if they: were not randomized, were primary prevention studies (ie. restricted to patients without

documented CAD), evaluated single-modality interventions (such as exercise-only programs, yoga interventions, or telephone follow-up), tested interventions delivered to hospitalized patients rather than outpatients, did not include a “usual care” arm, tested interventions that were not provided by health professionals (such as letter reminders, self help groups, self-directed interventions, or general health promotion interventions), or reported outcome data from less than 50 subjects. Studies in which patients with multiple diseases were enrolled were included if the outcomes for patients with coronary heart disease were reported separately or that data was provided by the study principal investigator when contacted.

Two of the investigators (AC and FM) independently assigned each reported intervention to one of 3 *a priori* defined groups: (1) Comprehensive Cardiac Rehabilitation (which included exercise as well as group education and counseling sessions about coronary risk factor management), (2) Cardiac Rehabilitation without exercise component (programs which included group education and counseling sessions about coronary risk factor management, but no structured exercise component), or (3) Individual Counselling (programs, usually delivered by specially trained nurses, involving

individual education and counseling sessions and individual follow-up, either in person or by telephone, to encourage coronary risk factor optimization). It should be noted that patient education was a key component of all 3 types of interventions (see Table 1 for a fuller description of the program in each included trial).

Statistical analysis:

Analyses were performed using RevMan 4.2 (The Cochrane Collaboration 2004). Our primary outcome was all-cause mortality at 12 months. Secondary outcomes that were meta-analyzed were recurrent myocardial infarctions and hospitalizations (although we attempted to obtain data on all-cause hospitalizations wherever possible, for some trials we were only able to obtain cardiovascular hospitalizations even after contact with the primary study authors). We defined “hospitalization rate” as the number of patients in each trial arm who were hospitalized at least once (thus, each patient could only contribute one event to these analyses).

As the outcomes were relatively common, risk ratios were calculated and the I^2 statistic was used to assess for heterogeneity in each outcome of interest. Studies were combined using the DerSimonian and Laird random effects model and the Peto fixed

effects model- where there was significant inter-trial heterogeneity, the results of the random effects model are reported, where there was no evidence of heterogeneity, the fixed effects results are presented. Analyses were conducted for each of the three types of programs: comprehensive cardiac rehabilitation, group cardiac rehabilitation (without exercise component), and individual counseling. For the primary analysis, we used data from the 12 month follow-up or, when unavailable, from the follow-up period closest to 12 months. We also conducted analyses for all programs combined using the various follow-up periods reported (6, 12, 24, 36, 48, 56, 60, and 72 months).

We described, but did not meta-analyze, the following outcomes: effects on major cardiovascular risk factors (cholesterol, smoking, blood pressure), use of proven efficacious therapies, patient quality of life, and patient functional status or symptom scores. These were evaluated and categorized as: statistically significant benefit seen in the intervention arm versus the control arm; trend towards better outcomes in the intervention arm, but did not reach statistical significance; or, no appreciable difference between the intervention and control arms. In order to standardize the

reporting of results for non-dichotomous outcomes (such as change in cholesterol or blood pressure levels, quality of life, or functional status scores), we calculated standardized effect sizes by dividing the absolute difference between intervention and control arms by the standard deviation in the control arms. By convention, effect sizes <0.20 are considered trivially small, 0.50 moderate, and >0.80 large.

RESULTS

Study selection and evaluation:

Overall we identified 6,336 citations from electronic databases (n=6,207), reference lists (n=36), and the Centres for Medicare and Medicaid Services (n=93). We reviewed 231 full manuscripts for potential inclusion. We excluded 180 of these studies after detailed evaluation; the reasons for exclusion are detailed in Figure 1 and Appendix A (a full list of excluded studies is included in Appendix B).

Disagreement among the reviewers regarding eligibility of the studies occurred on 16 occasions for a kappa value of 0.81. All disagreements were resolved by consensus.

Of the randomized trials eligible for inclusion,[28-79] 10 were reported in more than one publication. Two trials reported different endpoints in two separate publications.[28,29,57,58] One trial[30] reported the outcomes for all patients enrolled (only 56% of whom had cardiac disease) and, in a separate publication[31], provided details of event rates in the subgroup of patients with cardiac disease. The WHO Trial[32] included 24 collaborating centres; however, the original investigators excluded 7 sites because of poor subject follow-up and 4 sites due to significant differences at baseline

between the intervention and control arms. We included the 3-year outcome data from the remaining 13 sites as one trial for the purposes of this analysis, an approach validated by the non-significant tests for statistical heterogeneity for all-cause mortality ($Q=15.7$, 11 df, $p=0.16$) and MI ($Q=15.9$, 11 df, $p=0.15$) and the fact that the summary risk ratios for both endpoints were identical under the random and fixed effects models. While the two Finnish centres in the WHO Trial published their results separately (and for multiple follow-up periods), we included only their 3-year outcome data with the other 11 WHO sites for consistency of data presentation.[33-35] In five cases, we identified studies that reported longer follow-up data from another relevant trial.[36-40]

Studies included in the systematic review:

Summary data from the 41 unique randomized trials eligible for this systematic review are presented in Table 1.[28-79] In all of the trials, patients randomized to the control groups received usual care (this was generally undefined). One trial[41] is presented twice in the Table 1 because it had two intervention groups (comprehensive cardiac rehabilitation and group counseling) as well as a usual care control arm.

Our search retrieved 29 trials not included in our previous systematic review (which was limited to the pre-1999 literature)[25] and 22 trials not included in a more recent systematic review of cardiac rehabilitation (which was limited to the pre-2003 literature)[22].

Quantitative data synthesis:

All-cause mortality: Only one of the 27 trials reported a statistically significant survival benefit with the intervention (Table 2, Figures 2-4). The summary RR for all 27 trials reporting this outcome (13 625 patients, random effects RR 0.89 [95% CI 0.79-1.01]; fixed effects RR 0.89 [95% CI 0.79-1.01]) confirms that these interventions have not yet been shown to improve survival at 12 months. There was no significant statistical heterogeneity between trials ($p=0.82$, $I^2=0\%$) and the 12 month all-cause mortality rate in the control patients was 7%.

Although there was no appreciable difference in the treatment effects with any of the 3 types of secondary prevention programs (Table 2, Figures 2-4), there were differences in effect over time- while the RR for all-cause mortality was 0.97 (95% CI 0.82-1.14) in the 18 trials (9192 patients, p for heterogeneity=0.90, $I^2=0\%$)

reporting 12 month outcome data, the RR for all-cause mortality was 0.53 (95% CI 0.31-0.92) in the 4 trials (1367 patients, p for heterogeneity=0.44, I-squared=0%)[40,41,43,77] reporting 24 month outcome data. The 24 month all-cause mortality rate in the control patients was 5.5% and the number needed to treat to prevent one death in the next 24 months was 39. Furthermore, pooling the data from the 4 trials (2225 patients)[28,40,42,39] reporting follow-up data from at least 5 years after initiation of the intervention program demonstrates these programs had a sustained beneficial effect: the RR for all-cause mortality was 0.76 (95% CI 0.62-0.92) at 5 years with no appreciable heterogeneity between the trials (p =0.94, I-squared=0%).

Re-infarction Rate: One of the 16 trials reporting this endpoint (Table 2, Figures 5-7) detected a significant difference between intervention and control patients and the summary RR for 12 month re-infarction rate for all 9210 patients was 0.89 (95% CI 0.77-1.04). There was no significant statistical heterogeneity between trials (p =0.31, I-squared=13%) and the 12 month re-infarction rate in the control patients was 7%. There was no appreciable difference in the treatment effects with any of the 3 types of secondary prevention

programs (Table 2, Figures 5-7), nor were there any differences in effect over time.

Hospitalization Rate: Two of the 13 trials (5751 patients) reporting hospitalization rates detected a significant difference between intervention and control patients and the summary random effects RR for hospitalization rates for all 5751 patients was 0.85 (95% CI 0.78-0.93)- see Figure 8. Although there was no significant statistical heterogeneity between trials ($p=0.24$, $I^2=20\%$), we report this result using the random effects model as some trials reported data on all-cause hospitalizations and some only cardiovascular hospitalizations. The hospitalization rate in the control patients was 33% and the number needed to treat to prevent one hospitalization was 21 (median length of follow-up in these trials was approximately 12 months).

Restricting our analysis to the 9 trials (3653 patients) which reported all-cause hospitalization rates reveals a summary random effects RR of 0.84 (95% CI 0.74-0.97). Restricting our analysis to the 7 trials (3233 patients) which reported cardiovascular hospitalization rates reveals a summary random effects RR of 0.76 (95% CI 0.58-0.98).

Sensitivity Analyses: Analyses failed to reveal any effect of publication year on the observed results (data not shown).

Publication Bias: There was no evidence of publication bias (see Funnel Plot in Figure 9). The results of Begg's Test ($p=0.62$) and Egger's Test ($p=0.61$) confirm this.

Processes of Care: Twenty-four trials tested the impact of these disease management programs on cardiovascular risk factors, with 20 demonstrating better cholesterol profiles in patients randomized to the interventions, although the differences were statistically significant in only 12 trials and the effect sizes were generally small to moderate (Table 3). Of the 20 trials that assessed the use of proven efficacious medications, 8 demonstrated statistically significantly better application of at least one of these therapies in the intervention patients, 2 demonstrated better prescribing in intervention patients but did not achieve statistical significance, and 10 failed to demonstrate any appreciable difference between intervention and control patients (Table 3). It should be noted that in many cases the failure to demonstrate improved processes of care with the intervention was because of improved risk factor management in control patients- for example, in one study that

followed patients for over 4 years, 55% of controls had been exposed to comprehensive secondary prevention clinics by the close of the study.[37]

Other Endpoints: Fourteen of the 26 trials evaluating quality of life or functional status reported statistically significantly better scores in those patients exposed to the intervention programs, although the effect sizes were generally small (Table 3). Only 6 of these trials[30,44,49,50,65,66] described the costs of the intervention- while 2[30,50] reported that their intervention was cost-saving, none performed formal cost-effectiveness analyses. Another trial did not report costs, but did report that patients in the intervention arm had less visits to physicians as outpatients, less emergency room visits, less laboratory testing, and less total hospital days in follow-up than control patients.[78] Another trial reported statistically significantly lower inpatient bed days in intervention arm patients over 4 years of follow-up compared to controls (Dr. M. Vale, personal communication, January 10 2005).[76]

DISCUSSION

In summary, the weight of the published randomized trial evidence suggests that comprehensive secondary prevention programs positively impact on processes of care (risk factor profiles, use of proven efficacious therapies) which are closely linked to subsequent morbidity and mortality in patients with CAD.[80] Pooling the data from those trials which reported subsequent rates of MI does reveal a trend towards an 11% reduction in recurrent MIs over a median follow-up of 12 months which is not statistically significant; the majority of these programs also demonstrate improved symptom scores, exercise tolerance, or quality of life in participants. The mortality benefit derived from participation in the secondary prevention programs we identified (47% at 2 years [number needed to treat to prevent one death in 24 months=39] and 24% at 5 years) became apparent with longer lengths of follow-up, a fact which is not surprising considering the natural history of atherosclerotic CAD (ie. changes in coronary risk factors would not be expected to produce immediate improvements in atherosclerotic plaque stability or coronary artery diameter). There was a statistically significant 15% reduction in hospitalizations (driven by a statistically significant 24%

reduction in cardiovascular hospitalizations) over a median follow-up of 12 months (number needed to treat to prevent one hospitalization=21). These early beneficial effects on hospitalizations mirror the findings of a recent systematic review of multidisciplinary strategies for patients with heart failure which found that such interventions reduce hospitalizations by 25% within 6 months of implementation.[81]

Although some comprehensive lifestyle modification programs under consideration by the Medicare Coverage Advisory Committee, including the Ornish Program, were not included in our analysis (because the relevant randomized trial evidence reported data on less than 50 patients), our analyses extend the evidence base from these other programs to prove significant benefits on hard clinical endpoints. The Lifestyle Heart Trial evaluated the Ornish Program in 48 patients with CAD and demonstrated that a 12 month program emphasizing a low fat vegetarian diet, smoking cessation, stress management, and moderate exercise with group psychosocial support and counseling sessions improved coronary risk factors and significantly decreased the frequency and severity of angina in intervention patients vs. controls.[82] Furthermore, quantitative

coronary angiography demonstrated regression of coronary atherosclerosis in 82% of intervention patients versus 42% of control patients ($p=0.009$).[82] It is worth noting that the control patients in the Lifestyle Heart Trial were following a Step II diet of the National Cholesterol Education Project. Subsequent studies have confirmed that this multi-component cardiac rehabilitation program can be successfully taught and implemented at various American sites[83] and would likely be cost-saving.[84] However, it should be noted that while this economic analysis suggested that care cost reductions in the order of 30% to 60% within the first year were possible, the analyses are based on observational data (two concurrent cohorts followed for one year in one study, matched claims data analyses in another study, and two studies comparing actual costs after participation in the Ornish Program versus predicted costs) rather than randomized trial evidence.[84] Of course, a one year time horizon is likely too short for fully evaluating the cost-effectiveness of secondary prevention programs and studies with 5 and 10 year time horizons are needed (and ongoing).

Previously published systematic reviews of cardiac rehabilitation in survivors of myocardial infarction have reported

survival benefits in the order of 20-24%. However, most of the trials included in those overviews evaluated single modality exercise-based interventions (and thus were not included in our overview). For example, a recently published meta-analysis reported a statistically significant 20% reduction in all-cause mortality in 8432 patients; however, closer inspection of their report reveals that 40% of the data in their mortality analysis came from 13 trials which evaluated exercise-only programs and 2 trials which were excluded from our systematic review because of lack of a usual care control arm.[22]

As activity levels are inversely proportional to cardiovascular mortality and exercise training confers substantial physiologic and clinical benefits (including changes in endothelial function, autonomic tone, inflammatory markers),[23,85] it is not surprising that those trials found greater treatment effects than the trials included in our review (which evaluated multidisciplinary secondary prevention interventions that were not primarily exercise-based). However, the choice of programs to evaluate in our review was driven by current guideline recommendations that secondary prevention programs should not be restricted to supervised exercise programs alone.[26,27] Our systematic review is the first to prove that secondary prevention

programs which are delivered by health care providers and are not restricted to supervised exercise training do provide tangible reductions in clinically relevant endpoints such as hospitalization and death in addition to their well documented beneficial effects on patient risk factor profiles and quality of life/functional status.

Why didn't the trials reporting 12 month outcome data (including over 9000 subjects) demonstrate a statistically significant survival benefit? First, 12 months was clearly too short to show a clear impact on mortality- a fact supported by the known pathophysiology of atherosclerotic CAD and by the data demonstrating a significant survival benefit in those studies reporting outcomes over 2 years or more. It should be emphasized that studies which did evaluate coronary angiographic lesions at baseline and after 12 months did report statistically significant regression rates in patients compliant with comprehensive lifestyle modifications within 12 months even without significant changes in metabolic profiles or medication usage.[39,82] Second, the patients included in these studies were at sufficiently low risk over the first year after enrollment that the likelihood of detecting a beneficial effect was remote- indeed, the control event rates in these trials were substantially lower than

those in other trials enrolling patients with clinically overt CAD. Third, the incremental benefit of secondary prevention programs over usual care may be very small in the settings in which the trials were carried out (where management in the “usual care” arm may be close to optimal already). Indeed, secondary prevention programs are likely to be most beneficial in those settings where usual care is sub-optimal. Finally, it is possible that the labelling of patients with one disease for special attention in a disease-specific management program may have led to sub-optimal care for their co-morbid conditions and, as a result, no real difference in all-cause mortality.[86]

The overall effectiveness of secondary prevention programs should be interpreted in the context of the unexplained inconsistencies in effectiveness between different types of programs. As the programs were evaluated as single entities, it was not possible to ‘open the black box’ to identify what the key characteristics of successful programs were, which components were most influential, or how particular program or setting characteristics influenced outcomes. The particular mechanisms of effect of interventions remain poorly understood. Translation of the theoretical benefits of

secondary prevention programs into real-world patient benefit is also dependent on suitable patients being referred to, accessing, and completing the programs. Health outcomes research has consistently demonstrated that, even in publicly funded health care systems where access is free, only a minority of patients (less than one quarter to one half) ultimately access these programs.[10,87,88] Moreover, those groups that are less likely to be referred, attend, and complete programs are often those in greatest need of additional support and risk reduction, such as women, the elderly, low income groups, and ethnic minorities.[87] These groups were underrepresented in the studies reviewed.

Generalizability of the trial data:

As can be seen from Table 1, these trials enrolled relatively young patients- some even excluded patients over the age of 65. This raises potential concerns about the generalizability of our findings to this increasingly large population that is especially vulnerable to CAD. However, there is evidence that elderly patients demonstrate similar benefits after secondary prevention programs as younger people.[89-91] For instance, exercise training of elderly patients can provide a significant improvement in exercise tolerance similar to that

experienced in younger individuals.[92] While it is now less common for programs to have age-based restrictions for entry, older people are frequently excluded from programs due to a lack of program capacity to address their complex health needs or limited resources.[93,94] The effectiveness of programs for older patients may therefore be dependent not only on program content but also on program capacity to provide effective care to older patients who frequently have multiple co-morbidities.

Women were also underrepresented in the studies reviewed and data was not available to examine results by gender. This imbalance is significant because cardiac disease remains the leading cause of death for women in most of the developed world,[95] but is often viewed erroneously as principally being a “disease of men”. Sex differences in the investigation and management of CAD have been evident for many years.[96] Consequently, the need for improved and more responsive management of CAD in women has now been recognized by international guidelines.[96] While there is no evidence of any gender-based barriers to program benefit, women are consistently identified as being less likely to access programs.[97] To increase the strength of evidence supporting the

benefits of programs to women, more women should be included in study samples and data presented that examines the effectiveness of programs in males versus females.

Finally, as with any intervention proven efficacious in trial settings, the applicability of this evidence to the “real-world setting”, where compliance is likely to be highly variable and generally lower than that observed in trial participants, is a potential concern. While this may lead some to conclude that the results we report should be viewed as a “best case scenario” for the impact of secondary prevention programs, we disagree as we believe this view neglects that fact that randomized trial participants assigned to the control arm also receive care which is better than usual care. Indeed, as we pointed out earlier, the incremental benefit of secondary prevention programs over usual care may be very small in the settings in which these trials were carried out, where management in the “usual care” arm was often close to optimal already. Indeed, it seems likely that secondary prevention programs will be more beneficial in other settings (perhaps more akin to the “real world” of current clinical practice) where usual care is sub-optimal.

Limitations of this Review:

As with all systematic reviews, this study has a number of potential limitations. The most obvious (the lack of blinding in outcome ascertainment, lack of detail on whether randomization was conducted properly or whether allocation concealment was achieved, and our inability to identify unpublished studies- although we did not find any evidence for publication bias) arise from the primary data and, as all tend to result in over-estimation of any treatment effects,[98] these factors should be taken into account when interpreting our summary estimates. Our interpretation of these trials and the generalizability of the programs described is hampered by the imprecise descriptions of the interventions and the lack of data to determine the incremental benefits of the various components of each intervention. While some may criticize our choice of primary endpoints as being too broad to detect differences in “cardiac” morbidity and mortality, we believe that it is most appropriate to look at all-cause mortality or hospitalization as interventions to reduce resource use in one area can have unanticipated effects in another. Finally, we are unable to make a definitive comment on the cost-

effectiveness and economic impact of the programs tested in these trials due to the paucity of data.

In summary, although there was substantial variability in the interventions offered and the studies enrolled highly selected populations, secondary prevention programs do improve processes of care, coronary risk factor profiles, and functional status/quality of life. While the optimal mix of interventions, including their frequency and duration, are unclear, these programs do reduce hospitalizations and long-term mortality in patients with known CAD. While these programs appear to reduce health care resource use, their cost-effectiveness has been inadequately evaluated thus far in the literature. Thus, we believe that any policy decisions to implement secondary prevention programs on a wide scale should be accompanied by plans to rigorously evaluate long-term clinical and economic outcomes in participants and non-participants.

ACKNOWLEDGEMENTS: The authors gratefully acknowledge the help of Dr. N.C. Campbell (references 28 and 29), Dr. R. West (reference 46), Dr. M. Naylor (references 30 and 50), Dr. N. Marchionni (reference 66), Drs. Vale and Sundararajan (reference 76), Dr. Otterstad and Ms. Peersen (reference 77), and Dr. W. Young (reference 78) in providing further details about their studies. The authors also thank the staff of the University of Alberta Evidence-based Practice Center who participated in generating this report: Carol Friesen (searching), John Russell (retrieving articles, administrative tasks), Marilyn Josefsson (retrieving articles, administrative tasks), Natasha Wiebe (screening), Michelle Tubman (screening), Kenneth Bond (kappa calculations).

REFERENCES

1. Fox CS, Evans JC, Larson MG, Kannel WB, Levy D. Temporal trends in coronary heart disease mortality and sudden cardiac death from 1950 to 1999. The Framingham Heart Study. *Circulation* 2004;110:522-7.
2. American Heart Association. Heart Disease and Stroke Statistics — 2005 Update. Dallas, Tex.: American Heart Association; 2004
3. Yusuf S, Reddy S, Ounpuu S, Anand S. Global burden of cardiovascular diseases, part I: General considerations, the epidemiologic transition, risk factors, and impact of urbanization. *Circulation* 2001;104:2746-53.
4. Yusuf S, Hawken S, Ounpuu S, Dans T, Avezum A, Lanas F, et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* 2004;364:937-52.
5. Majumdar SR, Tsuyuki RT, McAlister FA. Controlling the ischemic heart disease epidemic: strategies for the 21st century. *Can J Public Health* 2001;92:9-11.
6. Yusuf S. Two decades of progress in preventing vascular disease. *Lancet* 2002;360:2-3.
7. Majumdar SR, McAlister FA, Furberg CD. From publication to practice in chronic cardiovascular disease – The Long and Winding Road. *J Am Coll Cardiol* 2004;43:1738-42.

8. EUROASPIRE I and II Group. Clinical reality of coronary prevention guidelines: a comparison of EUROASPIRE I and II in nine countries. Lancet 2001;357:995-1001.
9. Stafford RS, Radley DC. The underutilization of cardiac medications of proven benefit, 1990 to 2002. J Am Coll Cardiol 2003;41:56-61.
10. Witt BJ, Jacobsen SJ, Weston SA, Killian JM, Meverden RA, Allison TG, et al. Cardiac rehabilitation after myocardial infarction in the community. J Am Coll Cardiol 2004;44:988-96.
11. McCormick D, Gurwitz JH, Lessard D, Yarzebski J, Gore JM, Goldberg RJ. Use of aspirin, beta-blockers, and lipid-lowering medications before recurrent acute myocardial infarction: missed opportunities for prevention? Arch Intern Med. 1999;159:561-67.
12. Majumdar SR, Gurwitz JH, Soumerai SB. Undertreatment of hyperlipidemia in the secondary prevention of coronary artery disease. J Gen Intern Med. 1999;14:711-17.
13. Campbell NC, Thain J, Dean HG, et al. Secondary prevention in coronary heart disease: baseline survey of provision in general practice. BMJ 1998;316:1430-4.
14. Brady AJB, Oliver MA, Pittard JB. Secondary prevention in 24 431 patients with coronary heart disease: survey in primary care. BMJ 2001;322:1463.

15. Califf RM, DeLong ER, Ostbye T, et al. Underuse of aspirin in a referral population with documented coronary artery disease. *Am J Cardiol* 2002;89:653-61.
16. World Health Organisation. *Adherence to long-term therapies: evidence for action*. Geneva: WHO, 2003.
17. Benner JS, Glynn RJ, Mogun H, et al. Long-term persistence in use of statin therapy in elderly patients. *JAMA* 2002;288:455-61.
18. O'Connor GT, Buring JE, Yusuf S, Goldhaber SZ, Olmstead EM, Paffenbarger RS, et al. An overview of randomized trials of rehabilitation with exercise after myocardial infarction. *Circulation* 1989;80:234-44.
19. Oldridge NB, Guyatt GH, Fischer ME, Rimm AA. Cardiac rehabilitation after myocardial infarction. Combined experience of randomized clinical trials. *JAMA* 1988;260:945-50.
20. Dinnes J, Kleijnen J, Leitner M, Thompson D. Cardiac rehabilitation. *Qual Health Care* 1999;8:65-71.
21. Jolliffe JA, Rees K, Taylor RS, Thompson D, Oldridge N, Ebrahim S. Exercise-based rehabilitation for coronary heart disease. *The Cochrane Database of Systematic Reviews* 2001, Issue 1. Art. No.: CD001800. DOI: 10.1002/14651858.CD001800
22. Taylor RS, Brown A, Ebrahim S, Jolliffe J, Noorani H, Rees K, et al. Exercise-based rehabilitation for patients with coronary heart disease:

systematic review and meta-analysis of randomized controlled trials. *Am J Med* 2004;116:682-92.

23. Shephard RJ, Balady GJ. Exercise as cardiovascular therapy. *Circulation* 1999;99:963-72.
24. Hunter DJ, Fairfield G. Disease management. *BMJ* 1997;315:50-3.
25. McAlister FA, Lawson FME, Teo KK, Armstrong PW. Randomised trials of secondary prevention programs in coronary heart disease: systematic review. *BMJ* 2001;323:957-62.
26. Balady GJ, Ades PA, Comoss P, Limacher M, Pina IL, Southard D, et al. Core components of cardiac rehabilitation / secondary prevention programs. *Circulation* 2000;102:1069-1073.
27. ACC/AHA. Guidelines for the management of patients with acute myocardial infarction. *Circulation* 1999;100:1016-1030.
28. Campbell NC, Thain J, Deans HG, Ritchie LD, Rawles JG, Squair JL. Secondary prevention clinics for coronary heart disease: randomised trial of effect on health. *BMJ* 1998;316:1434-7.
29. Campbell NC, Ritchie LD, Thain J, Deans HG, Rawles JG, Squair JL. Secondary prevention in coronary heart disease: a randomised trial of nurse led clinics in primary care. *Heart* 1998;80:447-52.
30. Naylor MD, Brooten D, Campbell R, Jacobsen BS, Mezey MD, Pauly MV, et al. Comprehensive discharge planning and home follow-up of hospitalized elders. A randomized clinical trial. *JAMA* 1999;281:613-20.

31. Naylor MD, McCauley KM. The effects of a discharge planning and home follow-up intervention on elders hospitalized with common medical and surgical cardiac conditions. *J Cardiovasc Nurs* 1999;14:44-54.
32. World Health Organization Report 84: Rehabilitation and comprehensive secondary prevention after acute myocardial infarction. Copenhagen;1983:1-99.
33. Kallio V, Hamalainen H, Hakkila J, Luurila OJ. Reduction in sudden deaths by a multifactorial intervention programme after acute myocardial infarction. *Lancet* 1979;ii:1091-4.
34. Hamalainen H, Luurila OJ, Kallio V, Knuts L-R, Arstila M, Hakkila J. Long-term reduction in sudden deaths after a multifactorial intervention programme in patients with myocardial infarction: 10-year results of a controlled investigation. *Eur Heart J* 1989;10:55-62.
35. Hamalainen H, Luurila OJ, Kallio V, Knuts L-R. Reduction in sudden deaths and coronary mortality in myocardial infarction patients after rehabilitation. *Eur Heart J* 1995;16:1839-44.
36. Taylor CB, Miller NH, Smith PM, DeBusk RF. The effect of a home-based, case-managed, multifactorial risk-reduction program on reducing psychological distress in patients with cardiovascular disease. *Journal of Cardiopulmonary Rehabilitation* 1997;17:157-62.
37. Murchie P, Campbell NC, Ritchie LD, Simpson JA, Thain J. Secondary prevention clinics for coronary heart disease: four year follow up of a randomised controlled trial in primary care. *BMJ* 2003;326:84.

38. Murchie P, Campbell NC, Ritchie LD, Deans HG, Thain J. Effects of secondary prevention clinics on health status in patients with coronary heart disease: 4 year follow-up of a randomized trial in primary care. *Family Practice* 2004;21(5):567-74.
39. Niebauer J, Hambrecht R, Velich T, Hauer K, Marburger C, Kaulberer B, et al. Attenuated progression of coronary artery disease after 6 years of multifactorial risk intervention. *Circulation* 1997;96:2534-41.
40. Cupples ME, McKnight A. Five year follow up of patients at high cardiovascular risk who took part in randomised controlled trial of health promotion. *BMJ* 1999;319:687-8.
41. P.RE.COR. Group. Comparison of a rehabilitation programme, a counselling programme and usual care after an acute myocardial infarction: results of a long-term randomized trial. *European Heart Journal* 1991;12:612-6.
42. Vermeulen A, Lie KI, Durrer D. Effects of cardiac rehabilitation after myocardial infarction: Changes in coronary risk factors and long-term prognosis. *Am Heart J* 1983;105:798-801.
43. Haskell WL, Alderman EL, Fair JM, Maron DJ, Mackey SF, Superko R, et al. Effects of intensive multiple risk factor reduction on coronary atherosclerosis and clinical cardiac events in men and women with coronary artery disease. The Stanford Coronary Risk Intervention Project (SCRIP). *Circulation* 1994;89:975-90.

44. DeBusk RF, Houston Miller N, Superko R, Dennis CA, Thomas RJ, Lew HT, et al. A case-management system for coronary risk factor modification after acute myocardial infarction. *Ann Intern Med* 1994;120:721-9.
45. Cupples ME, McKnight A. Randomised controlled trial of health promotion in general practice for patient at high cardiovascular risk. *BMJ* 1994;309:993-6.
46. Jones DA, West RR. Psychological rehabilitation after myocardial infarction: multicentre randomized controlled trial. *BMJ* 1996;313:1517-21.
47. Frasure-Smith N, Lesperance F, Prince RH, Verrier P, Garber RA, Juneau M, et al. Randomised trial of home-based psychosocial nursing intervention for patients recovering from myocardial infarction. *Lancet* 1997;350:473-9.
48. Jolly K, Bradley F, Sharp S, Smith H, Thompson S, Kinmonth AL, et al for the SHIP Collaborative Group. Randomised controlled trial of follow up care in general practice of patients with myocardial infarction and angina: final results of the Southampton heart integrated care project (SHIP). *BMJ* 1999;318:706-11.
49. Fitzgerald JR, Smith DM, Martin DK, Freedman JA, Katz BP. A case manager intervention to reduce readmissions. *Arch Intern Med* 1994;154:1721-9.

50. Naylor M, Brooten D, Jones R, Lavizzo-Mourey R, Mezey M, Pauly M.
Comprehensive discharge planning for the hospitalized elderly. *Ann Intern Med* 1994;120:999-1006.
51. Allen JK, Blumenthal RS, Margolis S, Young DR, Miller ER III, Kelly K.
Nurse case management of hypercholesterolemia in patients with coronary heart disease: results of a randomized clinical trial. *American Heart Journal* 2002;144:678-86.
52. Allison TG, Farkouh ME, Smars PA, Evans RW, Squires RW, Gabriel SE, et al. Management of coronary risk factors by registered nurses versus usual care in patients with unstable angina pectoris (A chest pain evaluation in the emergency room (CHEER) substudy). *American Journal of Cardiology* 2000;86:133-8.
53. Allison TG, Squires RW, Johnson BD, Gau GT. Achieving national cholesterol education program goals for low-density lipoprotein cholesterol in cardiac patients: importance of diet, exercise, weight control, and drug therapy. *Mayo Clinic Proceedings* 1999;74:466-73.
54. Bengtsson K. Rehabilitation after myocardial infarction. A controlled study. *Scandinavian Journal of Rehabilitation Medicine* 1983;15:1-9.
55. Carlsson R. Serum cholesterol, lifestyle, working capacity and quality of life in patients with coronary artery disease. Experiences from a hospital-based secondary prevention programme. *Scandinavian Cardiovascular Journal Supplement* 1998;50:1-20.

56. Carlsson R, Lindberg G, Westin L, Israelsson B. Influence of coronary nursing management follow up on lifestyle after acute myocardial infarction. *Heart* 1997;77:256-9.
57. Engblom E, Ronnema T, Hamalainen H, Kallio V, Vanttinen E, Knuts L-R. Coronary heart disease risk factors before and after bypass surgery: results of a controlled trial on multifactorial rehabilitation. *European Heart Journal* 1992;13:232-7.
58. Engblom E, Korpilahti K, Hamalainen H, Ronnema T, Puukka P. Quality of life and return to work 5 years after coronary artery bypass surgery. Long-term results of cardiac rehabilitation. *J Cardiopulm Rehabil* 1997;17:29-36.
59. Fridlund B, Hogstedt B, Lidell E, Larsson PA. Recovery after myocardial infarction. Effects of a caring rehabilitation programme. *Scandinavian Journal of Caring Science* 1991;5(1):23-32.
60. Higgins HC, Hayes RL, McKenna KT. Rehabilitation outcomes following percutaneous coronary interventions (PCI). *Patient Education and Counseling* 2001;43:219-30.
61. Johnston M, Foulkes J, Johnston DW, Pollard B, Gudmundsdottir H. Impact on patients and partners of inpatient and extended cardiac counselling and rehabilitation: a controlled trial. *Psychosomatic Medicine* 1999;61:225-33.

62. Lear SA, Ignaszewski A, Linden W, Brozic A, Kiess M, Spinelli JJ, et al.
The extensive lifestyle management intervention (ELMI) following cardiac rehabilitation trial. *European Heart Journal* 2003;24:1920-7.
63. Lichtman JH, Amatruda J, Yaari S, Cheng S, Smith GL, Mattera JA, et al.
Clinical trial of an educational intervention to achieve recommended cholesterol levels in patients with coronary artery disease. *American Heart Journal* 2004;147:522-8.
64. Lisspers J, Sundin O, Hofman-Bang C, Nordlander R, Nygren A, Ryden L, et al.
Behavioral effects of a comprehensive, multifactorial program for lifestyle change after percutaneous transluminal coronary angioplasty: a prospective, randomized, controlled study. *Journal of Psychosomatic Research* 1999;46(2):143-54.
65. Masley S, Phillips S, Copeland JR. Group office visits change dietary habits of patients with coronary artery disease. The dietary intervention and evaluation trial (D.I.E.T.). *Journal of Family Practice* 2001;50(3):235-9.
66. Marchionni N, Fattirolli F, Fumagalli S, Oldridge N, Del Lungo F, Morosi L, et al.
Improved exercise tolerance and quality of life with cardiac rehabilitation of older patients after myocardial infarction. Results of a randomized, controlled trial. *Circulation* 2003;107:2201-6.
67. McHugh F, Lindsay GM, Hanlon P, Hutton I, Brown MR, Morrison C, et al.
Nurse led shared care for patients on the waiting list for coronary artery bypass surgery: a randomised controlled trial. *Heart* 2001;86:317-23.

68. Moher M, Yudkin P, Wright L, Turner R, Fuller A, Schofield T, et al for the Assessment of Implementation Strategies (ASSIST) Trial Collaborative Group. Cluster randomised controlled trial to compare three methods of promoting secondary prevention of coronary heart disease in primary care. *BMJ* 2001;322:1-7.
69. Oldridge N, Guyatt G, Jones N, Crowe J, Singer J, Feeny D, McKelvie R, et al. Effects on quality of life with comprehensive rehabilitation after acute myocardial infarction. *American Journal of Cardiology* 1991;67:1084-9.
70. Schuler G, Hambrecht R, Schlierf G, Niebauer J, Hauer K, Neumann J, et al. Regular physical exercise and low-fat diet. Effects on progression of coronary artery disease. *Circulation* 1992;86(1):1-11.
71. Sivarajan ES, Bruce RA, Lindskog BD, Almes MJ, Belanger L, Green B. Treadmill test responses to an early exercise program after myocardial infarction: a randomized study. *Circulation* 1982;65(7):1420-8.
72. Stagmo M, Westin L, Carlsson R, Israelsson B. Long-term effects on cholesterol levels and the utilization of lipid-lowering drugs of a hospital-based programme for secondary prevention of coronary artery disease. *Journal of Cardiovascular Risk* 2001;8:243-8.
73. Stern MJ, Gorman PA, Kaslow L. The group counselling v exercise therapy study. A controlled intervention with subjects following myocardial infarction. *Archives of Internal Medicine* 1983;143:1719-25.

74. Sundin O, Lisspers J, Hofman-Bang C, Nygren A, Ryden L, Ohman A.
Comparing multifactorial lifestyle interventions and stress management in coronary risk reduction. *International Journal of Behavioral Medicine* 2003;10(3):191-204.
75. Vale MJ, Jelinek MV, Best JD, Santamaria JD. Coaching patients with coronary heart disease to achieve the target cholesterol: a method to bridge the gap between evidence-based medicine and the “real world” – randomized controlled trial. *Journal of Clinical Epidemiology* 2002;55:245-52.
76. Vale MJ, Jelinek MV, Best JD, Dart AM, Grigg LE, Hare DL, et al.
Coaching patients on achieving cardiovascular health (COACH). *Archives of Internal Medicine* 2003;163:2775-83.
77. The Vestfold Heartcare Study Group. Influence on lifestyle measures and five-year coronary risk by a comprehensive lifestyle intervention programme in patients with coronary heart disease. *European Journal of Cardiovascular Prevention and Rehabilitation* 2003;10:429-37.
78. Young W, Rewa G, Goodman SG, Jaglal SB, Cash L, Lefkowitz C, et al.
Evaluation of a community-based inner-city disease management program for postmyocardial infarction patients: a randomized controlled trial. *CMAJ* 2003;169(9):905-10.
79. Yu C-M, Li LS-W, Ho HH, Lau C-P. Long-term changes in exercise capacity, quality of life, body anthropometry, and lipid profiles after a

cardiac rehabilitation program in obese patients with coronary heart disease. *American Journal of Cardiology* 2003;91(1):321-5.

80. Mant J, Hicks N. Detecting differences in quality of care: the sensitivity of measures of process and outcome in treating acute myocardial infarction. *BMJ* 1995;311:793-6.
81. McAlister FA, Stewart S, Ferrua S, McMurray JJV. Multidisciplinary strategies for the management of heart failure patients at high risk for admission: A systematic review of randomized trials. *J Am Coll Cardiol* 2004;44:810-9.
82. Ornish D, Brown SE, Scherwitz LW, Billings JH, Armstrong WT, Ports TA, et al. Can lifestyle changes reverse coronary heart disease? The Lifestyle Heart Trial. *Lancet* 1990;336:129-33.
83. Koertge J, Weidner G, Elliott-Eller M, Scherwitz L, Merritt-Worden TA, Marlin R, et al. Improvement in medical risk factors and quality of life in women and men with coronary artery disease in the Multicentre Lifestyle Demonstration Project. *Am J Cardiol* 2003;91:1316-22.
84. DeVries A, Palmer A, Scheib C, Day B, Bennett S, Fetterolf D, Kantz T. Highmark Blue Cross Blue Shield. Medical Management Analysis: Financial Impact of the Dr. Dean Ornish Program for Reversing Heart Disease. February 14, 2002;4:3.
85. Hambrecht R, Wolff A, Gielen S, et al. Effect of exercise on coronary endothelial function in patients with coronary artery disease. *N Engl J Med* 2000;342:454-60.

86. Redelmeier DA, Tan SH, Booth GL. The treatment of unrelated disorders in patients with chronic medical diseases. *N Engl J Med* 1998;338:1516-20.
87. Cooper AF, Jackson G, Weinman J, Horne R. Factors associated with cardiac rehabilitation attendance: a systematic review of the literature. *Clinical Rehabilitation* 2002;16:541-552.
88. Thomas RJ, Miller NH, Lamendola C, Berra K, Hedback B, Durstine JL, et al. National survey of gender differences in cardiac rehabilitation programs. Patient characteristics and enrollment patterns. *Journal of Cardiopulmonary Rehabilitation* 1996;16(6):402-12.
89. Milani RV, Lavie CJ. Prevalence and effects of cardiac rehabilitation on depression in the elderly with coronary heart disease. *American Journal of Cardiology* 1998;81(10):1233-6.
90. Majeed FA, Cook DG. Age and sex differences in the management of ischaemic heart disease. *Public Health* 1996;110:7-12.
91. Lavie CJ, Milani RV. Cardiac rehabilitation and preventive cardiology in the elderly. *Cardiology Clinics*. 1999;17(1):233-42.
92. Ades PA, Waldmann ML, Gillespie C. A controlled trial of exercise training in older coronary patients. *Journal of Gerontology* 1995;50A:M7-11.
93. Clark AM, Sharp C, MacIntyre PD. The role of age in moderating access to cardiac rehabilitation in Scotland. *Ageing and Society* 2002;22:501-515.
94. Oldridge N. Cardiac rehabilitation in the elderly. *Aging (Milano)* 1998;10(4):273-83.

95. British Heart Foundation. Coronary Heart Disease Statistics. London: BHF, 2004.
96. Mosca L, Appel LJ, Benjamin EJ, Berra K, Chandra-Strobos N, Fabunmi RP, et al. Evidence-Based Guidelines for Cardiovascular Disease Prevention in Women. *Circulation* 2004;109(5):672-693.
97. Cooper AF, Jackson G, Weinman J, Horne R. Factors associated with cardiac rehabilitation attendance: a systematic review of the literature. *Clinical Rehabilitation* 2002;16:541-552.
98. Schulz KF, Chalmers I, Hayes RJ, Altman DG. Empirical evidence of bias. *JAMA* 1995;273:408-12.

Don't have full report (but potentially relevant to the review, may be included in final report if we get data from the author before the mid-February final deadline):

Bell JA. Comparison of a multi-disciplinary home based cardiac rehabilitation programme with comprehensive conventional rehabilitation in post-myocardial infarction patients (PhD thesis). London, United Kingdom: University of London; 1998.

Box 1: Modifiable coronary risk factors (adapted from reference 4)

Modifiable Risk Factor	Prevalence in cases with myocardial infarction	Population Attributable Risk (99% CI)	Odds Ratio (99% CI) adjusted for age, gender, and smoking
Smoking	65%	36% (34% to 39%)	2.3 (2.1-2.4)
Dyslipidemia	33%	54% (50% to 59%)	3.9 (3.4-4.4)
Diabetes Mellitus	18%	12% (11% to 14%)	3.1 (2.8-3.4)
Hypertension	39%	23% (22% to 25%)	2.5 (2.3-2.7)
Abdominal Obesity	46%	34% (30% to 37%)	2.2 (2.1-2.5)
Psychosocial Factors	-	29% (23% to 36%)	2.5 (2.2-2.9)
Daily consumption of fruits and vegetables	36%	13% (10% to 17%)	0.7 (0.6-0.8)
Regular physical activity	14%	26% (20% to 32%)	0.7 (0.7-0.8)
Regular alcohol consumption	24%	14% (9% to 20%)	0.8 (0.7-0.9)
All of the above combined	-	90% (88% to 92%)	129.2 (90.2-185.0)

In the INTERHEART Study, “dyslipidemia” was defined as ApoB/ApoA1 Ratio in top quintile vs. lowest quintile; “abdominal obesity” was defined as waist/hip ratio > 0.90 in men and >0.83 in women; “psychosocial factors” was defined as positive exposure to depression, perceived stress at work or home, moderate or severe financial stress, low locus of control, and/or major life events; “regular physical activity” was defined as moderate or strenuous exercise for at least 4 hours per week; “regular alcohol consumption” was defined as 3 or more times per week.

Table 1: Description of studies included

Study	Sample Size	Study Population (Location)	Mean Age	% Male	Key Components of Intervention	Duration of Intervention
Comprehensive Cardiac Rehabilitation (15 trials, 3896 patients)						
Sivarajan (1982)	258 (170 in control and comprehensive secondary prevention arms)	Patients younger than 70 years discharged after AMI (USA)	57	>80%	Exercise program plus group education/counseling sessions about risk factor management	3 m
Vermeulen et al. (1983)	98	Males 40-55 yrs, discharged after AMI (Netherlands)	49	100%	Multidisciplinary team (details not given) involved in exercise rehabilitation, social and psychological supports for patients	1.5-2 m
Bengtsson (1983)	87	Patients aged <65 years, one year after AMI (Sweden)	56	85%	Rehabilitation program involving physical assessment and training by physiotherapy and counseling.	3m
World Health Organization* (1983)	1,735	Males < 65 yrs, discharged after AMI (Europe)	53	100%	Multidisciplinary team (components differed at each center) involved in patient health education and supervised exercise program	36 m
Oldridge (1991)	201	Patients discharged with diagnosis of AMI and evidence of anxiety or depression (Canada)	52	89%	Exercise prescription, supervised training and behavioral counseling	2 m
PRECOR (1991)	182	Males <65 years, discharged after AMI (France)	51	100%	Two intervention arms, one of which was: Comprehensive cardiac rehabilitation (supervised exercise program, relaxation training, risk factor management, education)	1.5 m
Fridlund et al. (1991)	178	Patients <65 years discharged after AMI (Sweden)	56	87%	Nurse-led rehabilitation program addressing lifestyle, stress and social support.	6
Engblom (1992)	228	Patients younger than 65 years, discharged after CABG (Finland)	54	88%	Group education, individual counseling (with physician and dietician) about diet and physical activity, supervised exercise training	0.75 m
Heidelberg Trial (Schuler 1992 and Niebauer 1997)	113	Males with CAD on angiography (Germany)	54	100%	Education about diet and exercise, exercise program with individual and group training sessions	12 m

Johnston et al. (1999)	100	Patients \leq 70 years hospitalized for 1 st time myocardial infarction (UK)	56	65%	Nurse-led inpatient and outpatient cardiac rehabilitation program containing education, support for risk factor change and psychological effects.	1.5 m
Lisspers et al (1999)	93	Patients <65 after PCI (Sweden)	53	37	Comprehensive residential (health education, behavioural change) containing skills training, habit rehearsal on stress management, smoking, diet, exercise and smoking; followed by outpatient program of self observation and reporting of risk factors with follow up support.	12m
Sundin et al. (2003)	132	Male patients <70 years after PCI, AMI or CABG (Sweden)	59	100%	Group-based multidisciplinary program addressing stress management, diet and exercise using lectures and skills training	12m
Yu et al. (2003)	112	Obese patients attending cardiac rehabilitation after acute MI or after percutaneous coronary intervention (China)	62	79%	Exercise program with group education classes about risk factor modification	2.5 m
Vestfold Heartcare Study (2003)	197	Patients discharged after acute coronary syndrome, CABG, PCI (85%); plus patients followed in clinic with stable CAD (15%) (Norway)	55	82%	Supervised exercise program, dietary advice, risk factor management education and individual plus group counseling involving a multidisciplinary team (physician, nurse, dietician, physiotherapist)	24 m
Marchionni et al (2003)	270	Patients older than 45 years discharged after AMI (Italy)	69	71%	Supervised exercise training and education/counseling about risk factor management, optional monthly support groups	2 m
Group Cardiac Rehabilitation without exercise component (4 trials, 2671 patients)						
Stern (1983)	106 (64 in control and group counseling arms)	Patients aged 30-69 years with recent MI (USA)	54	83%	Nurse and psychiatrist/social worker led group education and counseling sessions (12 sessions)	3 m
PRECOR (1991)	182	Males <65 years, discharged after AMI (France)	51	100%	Two intervention arms, one of which was: Group Counselling Program (group education and counseling led by physician, psychiatrist, and nutritionist)	1.5 m

Jones et al. (1996)	2328	Patients discharged home within 28 days of AMI (United Kingdom)	62	73%	Nurse and psychologist regularly saw participants for education, counselling, and relaxation/stress management training	1.75 m
DIET (2001)	97	Patients with known CAD and hyperlipidemia in specialty clinics (USA)	65	70%	Nurse-led education (group) and provision of written materials about diet and physical activity	12 m
Individual Counselling (23 trials, 11 896 patients)						
SCRIP (1994)	300	Patients < 75 yrs referred for angiography for known or suspected CAD (USA)	56	86%	Nurse-managed patient education and algorithm-driven management of risk factors, exercise program, frequent telephone and clinic visits with nurse	48 m
DeBusk et al. (1994) and Taylor (1997)	585	Patients ≤ 70 yrs discharged after AMI (USA)	57	79%	Nurse-managed patient education and counselling, exercise program, frequent telephone contact, and algorithm-based lipid therapy	12 m
Fitzgerald et al. (1994)	668	Patients > 45 yrs discharged from a general medicine in-patient service (2/3 with heart disease) and being followed at the general medicine clinic of a Veterans Affairs hospital (USA)	65	100%	Nurse-managed patient education, coordination of care, frequent telephone contact, and protocol-driven systematic assessments for unmet socio-medical needs	12 m
Naylor et al. (1994)	276 (142 with cardiac disease)	Patients > 70 yrs discharged from a tertiary care hospital with either CAD or heart failure (USA)	76	49%	Comprehensive discharge planning protocol with gerontologic nurse providing education, coordinating care, and maintaining telephone contact for 2 weeks	0.5 m
Cupples et al. (1994 and 99)	688	Patients <75 years with angina for at least 6 months identified from general practice records (UK)	63	59%	Individual nurse-led personalized health promotion program every 4 months	24 m
M-HART (1997)	1376	Patients discharged after AMI (Canada)	59	66%	Nurse contacted patients monthly by telephone, providing education and advice and screening patients for psychological distress- nurses referred patients to other health care resources as needed	12 m
Carlsson et al (1997)	168	Patients aged 50-70 years discharged after AMI (Sweden)	62	75%	Nurse-run education program (individual and group), exercise training program, nurse clinic visits	12 m
Carlsson et al	530	Patients aged 50-70 years discharged	62	79%	Individualized assessment and nurse	12 m

(1998)		after AMI, CABG or PCI (Sweden)			counseling on risk factors and diet	
Campbell et al. (1998), with longer term f/u reported in Murchie et al (2003)	1343	Patients <80 yrs old with documented CAD recruited from general practice records (United Kingdom)	66	58%	Regular followup at secondary prevention clinics run by nurses, promoting medical and lifestyle approaches to prevention	12 m
Jolly et al. (1999)	597	Patients with AMI or recent onset angina discharged from hospital or seen in a chest pain clinic (United Kingdom)	64	71%	Cardiac liaison nurse coordinated care between discharging service and family physician, patients given personal health record and prompts for follow-up	12 m
Naylor et al. (1999)	363 (202 with cardiac disease)	Patients ≥ 65 years discharged from a tertiary care hospital with either CAD or heart failure or after CABG/heart surgery (USA)	75	50%	Nurse-led patient education, coordination of home care, at least 2 home visits, use of a standardized protocol to optimize medications, and weekly telephone contact for 1 month	1 m
Allison et al. (1999)	152	Patients not treated with lipid lowering medication that completed cardiac rehabilitation after an acute coronary event (USA)	64	82%	Nurse-led follow up program every 6 weeks after start or change in lipid lowering therapy, including diet and exercise advice and lipid lowering medications.	6 m
Allison et al. (2000)	326	Patients attending emergency room with confirmed unstable angina (USA)	58	56%	Nurse-intervention including lipid management, referral to support services, counseling on risk factors and physician collaboration on abnormal results, 2 1-hour sessions at least 6 and 25 days after discharge	1 m
Moher et al. (2001)	1906	Patients 55-75 years identified in family practices with established CAD (UK)	66	68%	Nurse-led clinic providing support for risk factor change using electronic disease register and recall system	1 m
Stagmo et al. (2001)	241	Patients 50-69 years hospitalized in a CCU due to MI or previous CABG (Sweden)	62	78%	Hospital-based secondary prevention program	12 m
McHugh et al. (2001)	98	Patients on a waiting-list for elective CABG (UK)	62	76%	Shared nurse-led care program of monthly health education and motivational interviewing	7 m
Higgins et al	105	Patients discharged after PCI	48	90%	Nurse-led individualized education, risk	12 m

(2001)		(Australia)			factor goal setting and self-monitoring with telephone feedback, 3 home visits	
Allen et al. (2002)	228	Patients \leq 75 years discharged after CABG or PCI who had hypercholesterolemia (USA)	60	63%	Nurse practitioner case management in partnership with patients' primary provider (nurse-directed education and lifestyle modification advice, nurse clinic visits, nurse prescribed medications if necessary, f/u telephone calls)	12 m
COACH pilot (2002)	245	Patients \leq 75 years discharged after coronary revascularization procedure (Australia)	61	75%	Personal coaching by dietician via 5 telephone sessions and 5 mailings to achieve coronary risk factor targets (education, negotiated lifestyle plan, emphasis on followup with primary care provider and empowerment to ask for medication, repeated measurements)	6 m
COACH (2003)	792	Patients discharged from 6 hospitals after CABG, PCI, AMI, coronary angiography (Australia)	59	77%	Personal coaching (delivered by nurses or dieticians) via 5 telephone sessions and 5 mailings to achieve coronary risk factor targets (education, negotiated lifestyle plan, emphasis on followup with primary care provider and empowerment to ask for medication, repeated measurements)	6 m
ELMI Trial (2003)	302	Patients discharged from 2 tertiary-care cardiac rehabilitation programs (Canada)	64	83%	Personal coaching by case manager delivered via telephone and in-person counseling sessions; if suboptimal coronary risk factors at 6 months, treatment algorithms with cover letter from cardiologist mailed to primary care physicians	12 m
Young et al (2003)	146	Patients discharged home after AMI (Canada)	69	60%	Patient education, at least 6 home visits by nurse, nurse communication with primary care providers, and nurse-initiated referral for specialty care (based on standardized pathway)	2 m
REACH Trial (2004)	756	Patients aged 30-80 years discharged from tertiary care hospital with documented coronary disease (USA)	64	71%	Nurse-based education and counseling about cholesterol and target levels delivered via telephone (4 calls in 9 m) and mailed	12 m

					educational materials about a variety of secondary prevention manoeuvres	
--	--	--	--	--	---	--

* As outlined in text, the results for 13 of the 24 collaborating centers in the World Health Organization Trial are included here. Reasons for the exclusion of the other 11 centers are given in the text.

CHD=coronary heart disease; CHF=congestive heart failure; SCRIP= Standard Coronary Risk Intervention Project; COACH= Coaching patients On Achieving Cardiovascular Health Study; ELMI= Extensive Lifestyle Management Intervention ; REACH= Reinforcing Education About Cholesterol; CABG= Coronary Artery Bypass Grafting surgery ; PCI= Percutaneous coronary intervention; CCU= Coronary Care Unit; AMI= Acute Myocardial Infarction

Table 2: Impact of interventions on all-cause mortality and recurrent myocardial infarctions.

Study	Length of Follow-up	All-cause mortality (#events/total # patients)			Recurrent Myocardial Infarctions* (#events/total # patients)		
		Intervention Arm	Control Arm	Risk Ratio	Intervention Arm	Control Arm	Risk Ratio
Comprehensive Cardiac Rehabilitation							
Sivarajan	6 m	3/86	2/84	1.47 (0.25, 8.55)	NR	NR	NR
Vermeulen et al.	60 m	2/47	5/51	0.43 (0.09, 2.13)	4/47	9/51	0.48 (0.16, 1.46)
Bengtsson et al.	12 m	10/81	6/90	1.85 (0.70, 4.87)	2/81	4/90	0.56 (0.10, 2.95)
WHO	36 m	146/893	161/842	0.86 (0.70, 1.05)	150/893	139/842	1.02 (0.82, 1.26)
Oldridge et al.	12 m	3/99	4/102	0.77 (0.18, 3.36)	NR	NR	NR
PRECOR** - comprehensive rehabilitation arm	24 m	0/60	4/61	0.11 (0.01, 2.05)	4/60	6/61	0.68 (0.20, 2.28)
Fridlund et al.	12 m	9/87	14/91	0.67 (0.31, 1.47)	4/87	14/91	0.30 (0.10, 0.87)
Engblom et al.	12 m	12/119	13/109	0.85 (0.40, 1.77)	8/119	16/109	0.46 (0.20, 1.03)
Heidelberg Trial	12 m	2/56	1/57	2.04 (0.19, 21.82)	2/56	4/57	0.51 (0.10, 2.67)
	72 m	5/43	8/53	0.77 (0.27, 2.18)	3/43	4/53	0.92 (0.22, 3.91)
Lisspers et al	12 m	0/46	1/41	0.30 (0.01, 7.12)	NR	NR	NR
Vestfold Heartcare Study	24 m	2/98	1/99	2.02 (0.19, 21.92)	4/99	3/99	1.33 (0.72, 1.05)
Marchionni et al	12 m	7/180	3/90	1.17 (0.31, 4.41)	1/180	3/90	0.17 (0.02, 1.58)
Sub-Total:	12 trials	196/1852	215/1717	0.87 (0.73, 1.05)	177/1622	198/1490	0.86 (0.71, 1.04)
Group Cardiac Rehabilitation without exercise component							
Stern	12 m	0/35	1/29	0.28 (0.01, 6.57)	3/35	2/29	1.24 (0.22, 6.94)
PRECOR** -counselling arm	24 m	5/61	4/61	1.25 (0.35, 4.43)	4/61	6/61	0.67 (0.20, 2.25)
Jones et al.	6 m	53/1168	58/1160	0.91 (0.63, 1.31)	43/1168	48/1160	0.89 (0.59, 1.33)
	12 m	79/1168	84/1160	0.93 (0.69, 1.26)			
Sub-Total:	3 trials	84/1264	89/1250	0.94 (0.70, 1.25)	50/1264	56/1250	0.88 (0.61, 1.28)
Individual Counselling							
SCRIP	12 m	1/145	0/155	3.21 (0.13, 78.06)	5/145	0/155	11.75 (0.66, 210.69)
	24 m	1/145	2/155	0.53 (0.05, 5.83)	5/145	3/155	1.78 (0.43, 7.32)
	36 m	2/145	2/155	1.07 (0.15, 7.49)	5/145	6/155	0.89 (0.28, 2.86)
	48 m	3/145	3/155	1.07 (0.22, 5.21)	6/145	11/155	0.58 (0.22, 1.54)

**January 17, 2005 DRAFT FOR DISCUSSION AT JANUARY 25, 2005 MEDICARE COVERAGE ADVISORY
COMMITTEE MEETING**

58

DeBusk et al.	12 m	12/293	10/292	1.20 (0.52, 2.72)	10/293	20/292	0.50 (0.24, 1.05)
Fitzgerald et al.	12 m	35/333	35/335	1.01 (0.65, 1.57)	NR	NR	NR
Cupples et al.	24 m	13/342	29/346	0.45 (0.24, 0.86)	NR	NR	NR
	60 m	47/342	65/346	0.73 (0.52, 1.03)	NR	NR	NR
M-HART	12 m	38/692	27/684	1.39 (0.86, 2.25)	44/692	42/684	1.04 (0.69, 1.56)
Carlsson et al (1998)	12 m	2/118	2/117	0.99 (0.14, 6.92)	NR	NR	NR
Campbell et al.	12 m	22/673	25/670	0.88 (0.50, 1.54)	13/540	12/518	1.04 (0.48, 2.26)
	56 m	100/673	128/670	0.78 (0.61, 0.99)	100/673	125/670	0.80 (0.63, 1.01)
Jolly et al.	12 m	15/277	23/320	0.75 (0.40, 1.42)	NR	NR	NR
Allison et al. (2000)	6m	2/158	2/168	1.06 (0.15, 7.46)	0/158	1/168	0.35 (0.01, 8.63)
COACH pilot	6 m	0/121	2/124	0.20 (0.01, 4.22)	NR	NR	NR
COACH	6 m	4/398	4/394	0.99 (0.25, 3.93)	NR	NR	NR
	48 m	32/398	32/394		NR	NR	NR
ELMI	12 m	1/151	3/151	0.33 (0.04, 3.17)	NR	NR	NR
Young et al	14 m	8/71	11/75	0.77 (0.33, 1.80)	NR	NR	NR
Sub-Total:	13 trials	153/3772	173/3831	0.90 (0.73, 1.11)	72/1828	75/1817	0.95 (0.69, 1.30)
TOTAL	27 trials**	433/6888	477/6737	0.89 (0.79, 1.01)	299/4714	323/4496	0.89 (0.77, 1.04)

* Data for all trials except that of Campbell et al., DeBusk et al, and Allison et al. are for the combined endpoint of nonfatal and fatal myocardial infarction. The Campbell et al trial only collected data on nonfatal reinfarction rate and total mortality (they were unable to dissect out causes of mortality). The Allison et al and DeBusk et al trials collected data on nonfatal myocardial infarction.

** Note that for PRECOR, there were 2 intervention arms and 1 control arm

NR= not reported, but e-mail sent to PI requesting data if study published within past 10 years

Table 3: Impact of interventions on other endpoints

Study	Major Cardiovascular Risk Factors			Use of proven efficacious therapies	Patient Quality of Life	Patient functional status or symptom scores
	Cholesterol	Smoking	BP			
Comprehensive Cardiac Rehabilitation						
Sivarajan	0	0	0	0	0	0
Vermeulen et al.	++	–	0	0	0	+
Bengtsson	0	0	++	0	–	0
WHO*	++	–	++	++	0	–
Oldridge	0	0	0	0	+	–
PRECOR - comprehensive rehabilitation arm	0	–	0	0	0	++
Fridlund	0	–	0	0	++	++
Engblom	–	++	–	–	++	0
Heidelberg Trial -12 m f/u	++	–	0	–	0	0
-72 m f/u	–	–	0	–	0	0
Johnston	0	0	0	0	++	++
Lisspers et al.	0	++	0	0	0	0
Sundin et al.	+	0	0	0	0	0
Yu	0	0	0	0	+	+
Vestfold Heartcare Study	–	++	–	–	++	++
Marchionni et al	0	0	0	0	0	++
Group Cardiac Rehabilitation without exercise component						
Stern	0	0	0	0	++	0
PRECOR -counselling arm	0	–	0	0	0	++
Jones et al.	0	–	0	–	–	–
DIET	+	0	0	0	0	0
Individual Counselling						
SCRIP	++	++	+	++	0	0

**January 17, 2005 DRAFT FOR DISCUSSION AT JANUARY 25, 2005 MEDICARE COVERAGE ADVISORY
COMMITTEE MEETING**

60

DeBusk et al.	++	++	0	++	0	++
Fitzgerald et al	0	0	0	0	0	0
Naylor et al. (94)	0	0	0	0	–	–
Cupples et al.	+	+	+	++	0	++
M-HART	0	0	0	0	+	0
Carlsson et al (1997)	0	+	0	0	0	0
Carlsson et al (1998)	++	0	0	++	0	0
Campbell et al. -12m f/u -56 m year f/u	++ +	– –	++ +	++ –	++ –	++ –
Jolly et al.	+	–	+	–	–	–
Naylor et al. (99)	0	0	0	0	+	+
Allison et al. (99)	+	++	0	++	0	0
Allison et al. (2000)	+	–	–	–	0	0
Moher et al	++	++	++	–	–	0
Stagmo	+	0	0	+	0	0
McHugh	++	++	++	0	++	++
Higgins et al.	+	–	0	0	0	++
Allen et al.	++	0	0	+	0	0
COACH pilot	++	0	0	–	0	0
COACH	++	–	++	++	++	++
ELMI	–	–	–	0	–	–
Young et al	0	0	0	0	0	0
REACH	–	0	0	–	0	0

Use of proven efficacious therapies” encompasses both increased prescription rate by clinicians and/or increased compliance by patients.

++ Statistically significant benefit seen in the intervention arm vs. control arm.

+ Trend towards better outcomes in the intervention arm, but didn’t reach statistical significance.

– No appreciable difference between the intervention arm and control arm.

0 Not reported in study.

Comment [I1]:

Figure 1. Flow of trials through the selection process




Figure 2: All-cause mortality in trials evaluating comprehensive cardiac rehabilitation

Review: Secondary prevention programmes in coronary heart disease (Jan 14; 10:20)
Comparison: 02 All-Cause Mortality
Outcome: 09 Comprehensive Cardiac Rehabilitation




Figure 3: All-cause mortality in trials evaluating group cardiac rehabilitation without exercise component

Review: Secondary prevention programmes in coronary heart disease (Jan 14; 10:20)
Comparison: 02 All-Cause Mortality
Outcome: 10 Group Cardiac Rehabilitation (without exercise component)




Figure 4: All-cause mortality in trials evaluating individual counselling

Review: Secondary prevention programmes in coronary heart disease (Jan 14; 10:20)
Comparison: 02 All-Cause Mortality
Outcome: 11 Individual Counselling




Figure 5: Recurrent myocardial infarctions in trials evaluating comprehensive cardiac rehabilitation

Review: Secondary prevention programmes in coronary heart disease (Jan 14; 10:20)
Comparison: 03 Recurrent Myocardial Infarctions
Outcome: 01 Comprehensive Cardiac Rehabilitation




Figure 6: Recurrent myocardial infarctions in trials evaluating group cardiac rehabilitation without exercise component

Review: Secondary prevention programmes in coronary heart disease (Jan 14; 10:20)
Comparison: 03 Recurrent Myocardial Infarctions
Outcome: 02 Group Cardiac Rehabilitation (without exercise component)




Figure 7: Recurrent myocardial infarctions in trials evaluating individual counseling

Review: Secondary prevention programmes in coronary heart disease (Jan 14; 10:20)
Comparison: 03 Recurrent Myocardial Infarctions
Outcome: 03 Individual Counselling





Figure 8: Hospitalization rates in trials evaluating secondary prevention programs


Review: Secondary prevention programmes in coronary heart disease (Jan 14; 10:20)
Comparison: 06 Hospitalizations
Outcome: 04 All Programs



These data depict risk ratios for the number of patients requiring at least one hospitalization during follow-up. $RR < 1$ are consistent with less hospitalizations in the intervention arm; $RR > 1$ are associated with less hospitalizations in the control arm. Note that the data from some studies (SCRIP, Heidelberg, M-HART, Allison, and COACH) are “cardiovascular hospitalizations” while for the other studies it is “all-cause hospitalizations”. E-mails have been sent to authors of studies reporting only “cardiovascular hospitalizations” requesting data on “all-cause hospitalizations”. See text for results for “all-cause hospitalization” studies and for “cardiovascular hospitalizations” studies separately.

Figure 9: Funnel Plot for all-cause mortality data

Review: Secondary prevention programmes in coronary heart disease (Jan 14; 10:20)
Comparison: 02 All-Cause Mortality
Outcome: 12 All Programs



Appendix A. Search Strategies

MEDLINE - Ovid Version: rel9.1.0

Searched December 16, 2004

Results: 2527 unique records

1. exp "Case Management"/
2. exp "Comprehensive Health Care"/
3. exp "Disease Management"/
4. exp "Health Services Research"/
5. exp "Home Care Services"/
6. exp "Clinical Protocols"/
7. exp "Patient Care Planning"/
8. exp "Quality of Health Care"/
9. exp REHABILITATION/
10. (nurse adj led adj1 clinic\$).ti,ab.
11. (special\$ adj1 clinic\$).ti,ab.
12. or/1-11
13. exp "Myocardial Ischemia"/ or "Myocardial Ischemia\$".ti,ab.
14. RANDOMIZED CONTROLLED TRIAL.pt.
15. ANIMAL/ not HUMAN/
16. 14 not 15
17. 12 and 13 and 16
18. limit 17 to (english language and yr=1999 - 2005)
19. remove duplicates from 18

- The same search was conducted in **EBM Reviews - Cochrane Central Register of Controlled Trials - Ovid Version: rel9.1.0** (4th Quarter 2004) on December 16, 2004.
- There were 141 unique results.

PubMed

Searched December 16, 2004

Results: 50 unique records

The following search was conducted:

("Case Management"[MeSH] OR "Comprehensive Health Care"[MeSH] OR "Disease Management"[MeSH] OR "Health Services Research"[MeSH] OR "Home Care Services"[MeSH] OR "Clinical Protocols"[MeSH] OR "Clinical Protocols"[MeSH] OR "Patient Care Planning"[MeSH] OR "Quality of Health Care"[MeSH] OR "Rehabilitation"[MeSH]) AND ("Myocardial Ischemia"[MeSH] OR Myocardial Ischemia Field: Title/Abstract)

Limits: Publication Date from 2004/01/01 to 2004/12/17, Randomized Controlled Trial

Web of Science

Searched December 17, 2004

Results: 606 unique records

The **Cited Reference Search** feature was used to search for studies that cited the included studies from the original article.

EMBASE - Ovid Version: rel9.1.0

1988 to 2004 Week 51

Searched December 20, 2004

Results: 1313 unique records

1. exp "Patient Care"/
2. exp "Health Care"/
3. exp "Disease Management"/
4. exp "Health Services Research"/
5. exp "Home Care"/
6. exp "Clinical Protocol"/
7. exp "Health Care Quality"/
8. exp REHABILITATION/
9. (nurse adj led adj1 clinic\$.ti,ab.
10. (special\$ adj1 clinic\$.ti,ab.
11. or/1-10
12. exp "Heart Muscle Ischemia"/ or exp "Ischemic Heart Disease"/ or exp "Coronary Heart Disease"/ or "Myocardial Ischemia\$.ti,ab.
13. RANDOMIZED CONTROLLED TRIAL/
14. 11 and 12 and 13
15. limit 14 to english
16. limit 15 to human
17. remove duplicates from 16
18. limit 17 to yr=1999 - 2005

CINAHL (Cumulative Index to Nursing & Allied Health Literature) - Ovid Version: rel9.1.0

1982 to December Week 2 2004

Searched December 21, 2004

Results: 9 unique records

1. exp "Case Management"/
2. *Health Care Delivery/
3. exp "Disease Management"/
4. exp "Health Services Research"/
5. exp Home Health Care/
6. exp Protocols/
7. exp "Quality of Health Care"/
8. exp REHABILITATION/
9. (nurse adj led adj1 clinic\$).ti,ab.
10. (special\$ adj1 clinic\$).ti,ab.
11. or/1-10
12. exp "Myocardial Ischemia"/ or "Myocardial Ischemia\$".ti,ab.
13. clinical trial.pt.
14. animals/
15. 13 not 14
16. and/11-12,15
17. limit 16 to yr=1999-2005

SIGLE - FIZ Karlsruhe – Version Interhost 3000

Searched December 21, 2004

Results: 53

CORONARY OR MYOCARDIAL

AND

Health services, health administration, community care services

Appendix B: List of Excluded Studies and Reasons for Exclusion (full references at end of table)

Author year	Source	Reason for exclusion
Ades and Coello 2000	Database (WOS)	Guideline / review article
Akosah, Schaper, Havlik et al 2002	Database	Not randomized
Aldana, Whitmer, Greenlaw et al 2003	CMS	Not randomized
Ammerman, Keyserling, Atwood et al 2003	Database (Medline)	Primary prevention
Angerer, Siebert, Kothny et al 2000	CMS	Not randomized
Anonymous 1982	Database	Primary prevention
Ariyo, Haan, Tangen et al 2000	CMS	Not randomized
Arthur, Daniels, McKelvie 2000	Database (Medline)	Did not report primary outcomes
Barnard, Massey, Cherny et al 1983	CMS	Evaluated interventions which were not comprehensive disease management systems
Barnes, Trieber, Turner et al 1999	CMS	Population not CHD
Bartels, Gerdes, Babin-Ebell et al 2002	CMS	Guideline / review article
Beckie 1989	Database	Did not report primary outcomes
Bennett, Blackall, Clapham et al 1989	Database	Not randomized
Bentsson 1983	Database	Methodological flaw (patients excluded after randomization)
Berglund, Nilsson, Ericksson et al 2000	Database	Primary prevention
Berkman, Blumenthal, Burg et al 2003	Database (Medline)	Evaluated interventions which were not comprehensive disease management systems
Bethell and Mullee 1990	Reference list	Evaluated interventions which were not comprehensive disease management systems
Billings, Scherwitz, Sullivan et al	CMS	Guideline / review article
Bjarnason-Wehrens, Benesch, Bischoff et al 2003	Database (WOS)	Non-English
Blair, Bryant, Bocuzzi 1988	Database	Not randomized
Blumenthal, Jiang, Babyak et al 1997	CMS	Not randomized
Bogden, Koontz, Williamson et al 1997	Database	Did not report primary outcomes

Boulay and Prud'homme 2004	Database (WOS)	Not randomized
Bramlet, King, Young et al 1997	Database	Not randomized
Brown, Zhao, Chait et al 2001	CMS	Drug interventions
Burell	CMS	Guideline / review article
Cambien, Richard, Ducimetiere et al 1981	Database	Primary prevention
Campbell, Ritchie, Thain et al 1998	Database (Embase)	Protocol only
Cannon, Braunwald, McCabe et al 2004	CMS	Evaluated interventions which were not comprehensive disease management systems
Caracciolo, Davis, Sopko et al 1995	CMS	Not randomized
Carlson, Johnson, Franklin et al 2000	Database (WOS)	No usual care arm
Carney, Blumenthal, Stein et al 2001	CMS	Not randomized
Castillo-Richmond, Schneider, Alexander et al 2000	CMS	Evaluated interventions which were not comprehensive disease management systems
Chinaglia, Gaschino, Asteggiano et al 2002	Database	Not randomized
Clark, Bakhai, Lacey et al 2004	CMS	Not randomized
Coleman, Grothaus, Sandhu et al 1999	Database	Did not report the outcomes for patients with CHD separately or included <50% patients with CHD
Corti, Fuster, Fayad et al 2002	CMS	Drug interventions
Coull, Taylor, Elton et al 2004	Database (Medline)	Evaluated interventions which were not comprehensive disease management systems
Council on Clinical Cardiology and Council on Nutrition, Physical Activity and Metabolism 2003	CMS	Guideline / review article
Cummings, Hughes, Weaver et al 1990	Database	Did not report the outcomes for patients with CHD separately or included <50% patients with CHD
Cundiff 2002	CMS	Guideline / review article
DeBusk, Haskell, Miller et al 1985	Database	Evaluated interventions which were not comprehensive disease management systems
DeBusk, Miller, Parker et al 2004	Database (WOS)	Population not CHD
De Lorgeril, Salen, Martin et al 1999	CMS	Evaluated interventions which were not comprehensive disease

		management systems
Denollet and Brutsaert 2001	CMS	Not randomized
Detry, Vierendel, Vanbutsele et al 2001	Database (WOS)	Not randomized
DeVries, Palmer, Scheib et al 2002	CMS	Not randomized
DeVries, Day, Scott 2003	CMS	Not randomized
Diehl 1998	CMS	Not randomized
Dugan, Cohen	CMS	Guideline / review article
Dugmore, Tipson, Phillips et al 1999	Database (Medline)	Evaluated interventions which were not comprehensive disease management systems
Eaker, Sullivan, Kelly-Hayes et al 2004	CMS	Not randomized
Eddy 2000	CMS	Guideline / review article
Ellingsen, Hjermann, Abdelnoor et al 2003	Database (Medline)	Primary prevention
Elliott-Eller, Weidner, Pischke 2003	CMS	Abstract
Engblom, Korpilahti, Hamalainen et al 1997	Database	Did not report primary outcomes
Esposito, Giugliano, Nappo et al 2004	CMS	Drug interventions
Esselstyn 1999	CMS	Guideline / review article
Family Heart Study Group 1994	Database	Primary prevention
Fields, Walton, Schneider et al 2002	CMS	Protocol only
Flanagan, Cox, Paine et al 1999	Database	Not randomized
Frasure-Smith and Prince 1985	Database	Not randomized
Frasure-Smith, Lesperance, Gravel et al 2000	CMS	Not randomized
Friedman, Thoreson, Gill et al 1984	Database	Not randomized
Galatius, Gustafsson, Kistorp et al 2003	Database	Not randomized
Geil, Anderson, Gustafson 1995	CMS	Evaluated interventions which were not comprehensive disease management systems
George and Goldberg 2001	CMS	Guideline / review article
Ghoncheh and Smith 2004	CMS	Population not CHD
Gielen, Schuler, Hambrecht 2001	CMS	Guideline / review article
Gleason, Bourdet, Koehn et al 2002	CMS	<50 patients
Gould, Ornish, Scherwitz et al	CMS	<50 patients

1995		
Gould, Ornish, Kirkeeide et al 1992	CMS	Protocol only
Grimm for the MRFIT 1983	Database	Primary prevention
Grundy, Cleeman, Merz et al 2004	CMS	Guideline / review article
Gulati, Pandey, Arnsdorf et al 2003	CMS	Not randomized
Hakim, Curb, Petrovitch et al 1999	CMS	Evaluated interventions which were not comprehensive disease management systems
Hambrecht, Walther, Mobius-Winkler 2004	CMS	Evaluated interventions which were not comprehensive disease management systems
Harris, Record, Gipson et al 1998	Database	Not randomized
Hedback and Perk 1987	Database	Not randomized
Imperial Cancer Research Fund OXCHECK Study Group 1995	Database	Primary prevention
Jain, Uppal, Bhatnagar et al 1993	CMS	Evaluated interventions which were not comprehensive disease management systems
Jukema, Bruschke, van Boven et al 1995	CMS	Drug interventions
Kawachi, Sparrow, Vokonas et al 1994	CMS	Not randomized
Ketola, Makela, Klockars 2001	Database (WOS)	Primary prevention
Koertge, Weidner, Billings et al 2002	CMS	Abstract
Koertge, Weidner, Elliott-Eller et al 2003	CMS	Not randomized
Kornitzer, De Backer, Dramaix et al 1980	Database	Primary prevention
Krachler	Reference list	<50 patients
Kris-Etherton, Harris, Appel et al 2002	CMS	Guideline / review article
Lampert, Joska, Burg et al 2002	CMS	Not randomized
Lear, Ignaszewski, Linden et al 2002	Database (WOS)	Protocol only
Lesperance, Frasure-Smith, Talajic et al 2002	CMS	Not randomized
Lewin, Furze, Robinson et al 2002	Database (Medline)	No usual care arm
Lewis and Resnik 1967	Database	Did not report the outcomes for

		patients with CHD separately or included <50% patients with CHD
Liao, Ma, Dong et al 2003	Database (Embase)	Non-English
Lichtenstein and Van Horn 1998	CMS	Guideline / review article
Lindholm, Ekblom, Dash et al 1995	Database	Primary prevention
Maggioni 2000	Database (Embase)	Guideline / review article
Malach and Imperato 2004	CMS	Not randomized
Marra, Paolillo, Spadaccini et al 1985	Database	Evaluated interventions which were not comprehensive disease management systems
Marshall, Penckofer, Llewellyn 1986	Database	Not randomized
Matthews, Gump, Harris et al 2004	CMS	Not randomized
Meer 1999	Database (SIGLE)	Not randomized
Meland, Laerum, Ulvik 1997	Database	Primary prevention
Merritt, Scherwitz, Brown et al 1990	CMS	<50 patients
Merritt, Ornish, Scherwitz et al 1995a	CMS	Abstract
Merritt, Ornish, Scherwitz et al 1995b	CMS	Abstract
Merritt-Worden, Pettengill, Ornish 2003	CMS	Abstract
Miettinen, Pyorala, Olsson et al 1997	CMS	Drug interventions
Miller, Erlinger, Young et al 2002	CMS	<50 patients
Mittleman, Maclure, Sherwood et al 1995	CMS	Not randomized
National Cholesterol Education Program, National Institutes of Health	CMS	Guidelines / review article
Ness, Hughes, Elwood et al 2002	Database (Medline)	Evaluated interventions which were not comprehensive disease management systems
Nicholson, Sklar, Barnard et al 1999	CMS	Evaluated interventions which were not comprehensive disease management systems
Oldenburg, Martin, Greenwood 1995	Database	Did not report primary outcomes
Ornish, Scherwitz, Doody et al	CMS	<50 patients

1983		
Ornish, Brown, Scherwitz et al 1990	CMS	<50 patients
Ornish 1998	CMS	Not randomized
Ornish, Scherwitz, Billings et al 1998	Database / CMS	<50 patients
Ornish 2002	CMS	Guideline / review article
Ornish and Pettengill 2003	CMS	Abstract
Ornish 2004	CMS	Guideline / review article
Ornish (chapter 8)	CMS	Guideline / review article
Ornish and Hart (chapter 34)	CMS	Guideline / review article
Pater, Ditlef Jacobsen, Rollag et al 2000	Database (Embase)	Protocol only (no data presented)
Peiss, Kurlito, Rubenfire 1995	Database	Not randomized
Pettengill, Pearson, Pifalo et al 2002	CMS	Abstract
Pfisterer, Buser, Osswald et al 2003	CMS	Not randomized
Picard, Schwartz, Ahn et al 1989	Database	Evaluated interventions which were not comprehensive disease management systems
Pischke, Weidner, Billings J et al 2002	CMS	Abstract
Pitt, Waters, Brown et al 1999	CMS	Drug interventions
Pollock, Franklin, Balady et al 2000	CMS	Guideline / review article
Pozen, Stechmiller, Harris et al 1977	Database	Inpatient-based intervention
Prochaska, Johnson, Lee	CMS	Guideline / review article
Pyke, Wood, Kinmonth et al 1997	Database	Primary prevention
Rahe, Ward, Hayes 1979	Database	<50 patients
Rihal, Raco, Gersh et al 2003	CMS	Guideline / review article
Roderick, Ruddock, Hunt et al 1997	Database	Primary prevention
Roman, Gutierrez, Luksic et al 1983	Database	Evaluated interventions which were not comprehensive disease management systems
Rose, Heller, Pedoe et al 1980	Database	Primary prevention
Rubenstein, Kahn, Reinisch et al 1990	Database	Not randomized
Ruo, Rumsfeld, Hlatky et al 2003	CMS	Not randomized
Scandinavian Simvastatin Survival Study 1994	CMS	Drug interventions

Schectman, Wolff, Byrd et al 1996	Database	Did not report the outcomes for patients with CHD separately or included <50% patients with CHD
Schneider, Staggers, Alexander et al 1995	CMS	Evaluated interventions with were not comprehensive disease management systems
Sdringola, Nakagawa, Nakagawa et al 2003	CMS	Not randomized
Shaffer and Wexler 1995	Database	Not randomized
Shintani, Beckham, Brown et al 2001	CMS	Population not CHD
Simpson, Dixon, Bolli 2004	Database (WOS)	Not randomized
Sivarajan, Newton, Almes et al 1983	Database	Did not report primary outcomes
The South East London Screening Study Group 1977	Database	Primary prevention
Specchia, De Servi, Scire et al 1996	Reference list	No usual care arm
Stahle, Mattsson, Ryden et al 1999	Database (Medline)	Evaluated interventions with were not comprehensive disease management systems
Starkey, Michaelis, Lusignan 2000	Database	Not randomized
Stern and Cleary 1982	Database	Evaluated interventions with were not comprehensive disease management systems
Strandberg, Pitkala, Berglind et al 2001	Database (Embase)	Protocol only (no data presented)
Taddei, Galetta, Virdis et al 2000	CMS	Evaluated interventions with were not comprehensive disease management systems
Thoresen, Friedman, Gill et al 1982	Database	Not randomized
Townsend, Piper, Frank et al 1988	Database	Evaluated interventions with were not comprehensive disease management systems
Tu, Pashos, Naylor et al 1997	CMS	Not randomized
Vale, Jelinek, Best et al 2003	Database (Medline)	Summary of trial already included
Van Drenth, Hulscher, Mokkink et al 1997	Database	Not randomized
Vedin, Wilhelmsson, Tibblin et al 1976	Database	Not randomized
Von Birgelen, Hartmann, Mintz et al 2003	CMS	Not randomized

Wallner, Watzinger, Lindschinger et al 1999	Database (Medline)	<50 patients
Wasson, Gaudette, Whaley et al 1992	Database	Evaluated interventions which were not comprehensive disease management systems
Waters, Higginson, Gladstone et al 1994	CMS	Drug interventions
Weber, Barnard, Roy 1983	CMS	Population not CHD
Weidner, Pischke, Eller 2003	CMS	Abstract
Weinberger, Smith, Katz et al 1988	Database	Did not report the outcomes for patients with CHD separately or included <50% patients with CHD
Weingarten, Reidinger, Conner et al 1994	Database	Inpatient-based intervention
Weintraub, Clements, Crisco et al 2003	CMS	Not randomized
Williams, Paton, Siegler et al 2000	CMS	Not randomized
Woollard, Burke, Beilin et al (Journal of Cardiovascular Risk) 2003	Database (Medline)	<50 patients
Woollard, Burke, Beilin (Journal of Human Hypertension) 2003	Database (WOS)	<50 patients
Yu-Poth, Zhao, Etherton et al 1999	CMS	Guideline / review article

Full Citations for **Excluded Studies**

Ades PA, Coello CE. Effects of exercise and cardiac rehabilitation on cardiovascular outcomes. *Med Clin North Am* 2000 Jan;84(1):251-65, x-xi.

Akosah KO, Schaper AM, Havlik P, Barnhart S, Devine S. Improving care for patients with chronic heart failure in the community. The importance of a disease management program. *Chest* 2002;122:906-12.

Aldana SG, Whitmer WR, Greenlaw R, Avins AL, Salberg A, Barnhurst M, et al. Cardiovascular risk reductions associated with aggressive lifestyle modification and cardiac rehabilitation. *Heart Lung* 2003 Nov-Dec;32(6):374-82.

Ammerman AS, Keyserling TC, Atwood JR, Hosking JD, Zayed H, Krasny C. A randomized controlled trial of a public health nurse directed treatment program for rural patients with high blood cholesterol. *Prev Med* 2003 Mar;36(3):340-51.

Angerer P, Siebert U, Kothny W, Muhlbauer D, Mudra H, von Schacky C. Impact of social support, cynical hostility and anger expression on progression of coronary atherosclerosis. *J Am Coll Cardiol* 2000 Nov 15;36(6):1781-8.

Anonymous. The Roman coronary heart disease prevention program. Final results. *Giornale Italiano di Cardiologia* 1982;12:541-54.

Ariyo AA, Haan M, Tangen CM, Rutledge JC, Cushman M, Dobs A, et al. Depressive symptoms and risks of coronary heart disease and mortality in elderly Americans. Cardiovascular Health Study Collaborative Research Group. *Circulation* 2000 Oct 10;102(15):1773-9.

Arthur HM, Daniels C, McKelvie R, Hirsh J, Rush B. Effect of a preoperative intervention on preoperative and postoperative outcomes in low-risk patients awaiting elective coronary artery bypass graft surgery. A randomized, controlled trial. *Ann Intern Med* 2000 Aug 15;133(4):253-62.

Barnard RJ, Massey MR, Cherny S, O'Brien LT, Pritikin N. Long-term use of a high-complex-carbohydrate, high-fiber, low-fat diet and exercise in the treatment of NIDDM patients. *Diabetes Care* 1983 May-Jun;6(3):268-73.

Barnes VA, Treiber FA, Turner JR, Davis H, Strong WB. Acute effects of transcendental meditation on hemodynamic functioning in middle-aged adults. *Psychosom Med* 1999 Jul-Aug;61(4):525-31.

Bartels C, Gerdes A, Babin-Ebell J, Beyersdorf F, Boeken U, Doenst T, et al. Working Group on Extracorporeal Circulation and Mechanical Ventricular Assist Devices of the German Society for Thoracic and Cardiovascular Surgery.

Cardiopulmonary bypass: Evidence or experience based? J Thorac Cardiovasc Surg 2002 Jul;124(1):20-7.

Beckie T. A supportive-educative telephone program: Impact on knowledge and anxiety after coronary artery bypass graft surgery. Heart Lung 1989;18:46-55.

Bennett P, Blackall M, Clapham M, Little S, Player D, Williams K. South Birmingham Coronary Prevention Project: a district approach to the prevention of heart disease. Community Medicine 1989;11:90-6.

Bentsson K. Rehabilitation after myocardial infarction. A controlled study. Scan J Rehab Med 1983;15:1-9.

Berglund G, Nilsson P, Ericksson KF, et al. Long-term outcome of the Malmo preventive project: mortality and cardiovascular morbidity. J Intern Med 2000;247:19-29.

Berkman LF, Blumenthal J, Burg M, Carney RM, Catellier D, Cowan MJ, et al. Enhancing Recovery in Coronary Heart Disease Patients Investigators (ENRICHD). Effects of treating depression and low perceived social support on clinical events after myocardial infarction: the Enhancing Recovery in Coronary Heart Disease Patients (ENRICHD) Randomized Trial. JAMA 2003 Jun 18;289(23):3106-16.

Bethell HJ, Mullee MA. A controlled trial of community based coronary rehabilitation. Br Heart J 1990;64:370-5.

Billings JH, Scherwitz LW, Sullivan R, Sparler S, Ornish DM. Chapter 9: The lifestyle heart trial: comprehensive treatment and group support therapy. No further details available (reference provided by CMS).

Bjarnason-Wehrens B, Benesch L, Bischoff KO, Buran-Kilian B, Gysan D, Hollenstein U, et al. Effects of a Phase II Cardiac Rehabilitation Program Performed on an Outpatient Basis. Herz 2003 Aug; 28(5):404-412.

Blair TP, Bryant FJ, Bocuzzi S. Treatment of hypercholesterolemia by a clinical nurse using a stepped-care protocol in a non-volunteer population. Arch Intern Med 1988;148:1046-8.

Blumenthal JA, Jiang W, Babyak MA, Krantz DS, Frid DJ, Coleman RE, et al. Stress management and exercise training in cardiac patients with myocardial ischemia. Effects on prognosis and evaluation of mechanisms. Arch Intern Med 1997 Oct 27;157(19):2213-23.

Bogden PE, Koontz LM, Williamson P, Abbott RD. The physician and pharmacist team. An effective approach to cholesterol reduction. *J Gen Intern Med* 1997;12:158-64.

Boulay P, Prud'homme D. Health-Care Consumption and Recurrent Myocardial Infarction After 1 Year of Conventional Treatment Versus Short- and Long-Term Cardiac Rehabilitation. *Preventive Medicine* 2004 May; 38(5):586-593.

Bramlet DA, King H, Young L, Witt JR, Stoukides CA, Kaul AF. Management of hypercholesterolemia: practice patterns for primary care providers and cardiologists. *Am J Cardiol* 1997;80(8B):39H-44H.

Brown BG, Zhao XQ, Chait A, Fisher LD, Cheung MC, Morse JS, et al. Simvastatin and niacin, antioxidant vitamins, or the combination for the prevention of coronary disease. *N Engl J Med* 2001 Nov 29;345(22):1583-92.

Burell G. Behavioral medicine interventions in secondary prevention of coronary heart disease. In: K. Orth-Gomer & N. Schneiderman (Eds.), *Behavioral medicine approaches to cardiovascular disease prevention* (pp. 227-236).

Cambien F, Richard JL, Ducimetiere P, Warnet JM, Kahn J. The Paris Cardiovascular Risk Factor Prevention Trial. Effects of two years of intervention in a population of young men. *J Epid Comm Health* 1981;35:91-7.

Campbell NC, Ritchie LD, Thain J, Deans HG, Rawles JM, Squair JL. Secondary prevention in coronary heart disease: a randomised trial of nurse led clinics in primary care. *Heart* 1998 Nov;80(5):447-52.

Cannon CP, Braunwald E, McCabe CH, Rader DJ, Rouleau JL, Belder R, et al. Intensive versus moderate lipid lowering with statins after acute coronary syndromes. *N Engl J Med* 2004 Apr 8;350(15):1495-504.

Caracciolo EA, Davis KB, Sopko G, Kaiser GC, Corley SD, Schaff H, et al. Comparison of surgical and medical group survival in patients with left main coronary artery disease. Long-term CASS experience. *Circulation* 1995 May 1;91(9):2325-34.

Carlson JJ, Johnson JA, Franklin BA, Vanderlaan RL. Program Participation, Exercise Adherence, Cardiovascular Outcomes, and Program Cost of Traditional Versus Modified Cardiac Rehabilitation. *American Journal of Cardiology* 2000 Jul 1; 86(1):17-23.

Carney RM, Blumenthal JA, Stein PK, Watkins L, Catellier D, Berkman LF, et al. Depression, heart rate variability, and acute myocardial infarction. *Circulation* 2001 Oct 23;104(17):2024-8.

Castillo-Richmond A, Schneider RH, Alexander CN, Cook R, Myers H, Nidich S, et al. Effects of stress reduction on carotid atherosclerosis in hypertensive African Americans. *Stroke* 2000 Mar;31(3):568-73.

Chinaglia A, Gaschino G, Asteggiano R, Titta G, Gullstrand R, Bobbio M, et al. Impact of a nurse-based heart failure management program on hospitalization rate, functional status, quality of life, and medical costs. *Ital Heart J* 2002;3:532-8.

Clark MA, Bakhai A, Lacey MJ, Pelletier EM, Cohen DJ. Clinical and economic outcomes of percutaneous coronary interventions in the elderly: an analysis of medicare claims data. *Circulation* 2004 Jul 20;110(3):259-64.

Coleman EA, Grothaus LC, Sandhu N, Wagner EH. Chronic care clinics: a randomized controlled trial of a new model of primary care for frail older adults. *J Am Geriatr Soc* 1999;47:775-83.

Corti R, Fuster V, Fayad ZA, Worthley SG, Helft G, Smith D, et al. Lipid lowering by simvastatin induces regression of human atherosclerotic lesions: two years' follow-up by high-resolution noninvasive magnetic resonance imaging. *Circulation* 2002 Dec 3;106(23):2884-7.

Coull AJ, Taylor VH, Elton R, Murdoch PS, Hargreaves AD. A randomised controlled trial of senior Lay Health Mentoring in older people with ischaemic heart disease: The Braveheart Project. *Age Ageing*. 2004 Jul;33(4):348-54.

Council on Clinical Cardiology and Council on Nutrition, Physical Activity and Metabolism. Exercise and physical activity in the prevention and treatment of atherosclerotic cardiovascular disease: a statement from the Council on Clinical Cardiology (Subcommittee on Exercise, Rehabilitation, and Prevention) and the Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity). *Circulation* 2003 Jun 24;107(24):3109-16.

Cummings JE, Hughes SL, Weaver FM, et al. Cost-effectiveness of Veterans Administration hospital-based home care. *Arch Intern Med* 1990;150:1274-80.

Cundiff DK. Coronary artery bypass grafting (CABG): reassessing efficacy, safety, and cost. *MedGenMed* 2002 Jun 19;4(2):7.

DeBusk RF, Haskell WL, Miller NH, Berra K, Taylor CB, Berger WE, et al. Medically directed at-home rehabilitation soon after clinically uncomplicated acute myocardial infarction: a new model for patient care. *Am J Cardiol* 1985;55:251-7.

Debusk RF, Miller NH, Parker KM, Bandura A, Kraemer HC, Cher DJ, et al. Care Management for Low-Risk Patients With Heart Failure - a Randomized, Controlled Trial. *Annals of Internal Medicine* 2004 Oct 19; 141(8):606-613.

de Lorgeril M, Salen P, Martin JL, Monjaud I, Delaye J, Marmelle N. Mediterranean diet, traditional risk factors, and the rate of cardiovascular complications after myocardial infarction: final report of the Lyon Diet Heart Study. *Circulation* 1999 Feb 16;99(6):779-85.

Denollet J, Brutsaert DL. Reducing emotional distress improves prognosis in coronary heart disease: 9-year mortality in a clinical trial of rehabilitation. *Circulation* 2001 Oct 23;104(17):2018-23.

Detry JMR, Vierendeel IA, Vanbutsele RJ, Robert AR. Early Short-Term Intensive Cardiac Rehabilitation Induces Positive Results as Long as One Year After the Acute Coronary Event: a Prospective One-Year Controlled Study. *Journal of Cardiovascular Risk* 2001 Dec; 8(6):355-361.

DeVries A, Day B, Scott D. Financial impact of the Dr. Dean Ornish program for reversing heart disease. In: *Medical Management Analysis*; Highmark; June 2003.

DeVries A, Palmer A, Scheib C, Day B, Bennett S, Fetterolf D, et al. Financial impact of the Dr. Dean Ornish program for reversing heart disease. In: *Medical Management Analysis; Lifestyle Advantage from Highmark and Dr. Dean Ornish*; Highmark, Blue Cross Blue Shield; 2002 February 14;4(3).

Diehl HA. Coronary risk reduction through intensive community-based lifestyle intervention: the Coronary Health Improvement Project (CHIP) experience. *Am J Cardiol* 1998 Nov 26;82(10B):83T-87T.

Dugan E, Cohen SJ. Improving physicians' implementation of clinical practice guidelines: enhancing primary care practice. In Shumaker SA, Schron EB, Ockene JK, McBee WL (Eds.). *The handbook of health behavior change – second edition* (pp. 283-304) Springer Publishing Company.

Dugmore LD, Tipson RJ, Phillips MH, Flint EJ, Stentiford NH, Bone MF, et al. Changes in cardiorespiratory fitness, psychological wellbeing, quality of life, and vocational status following a 12 month cardiac exercise rehabilitation programme. *Heart* 1999;81:359-66.

Eaker ED, Sullivan LM, Kelly-Hayes M, D'Agostino RB Sr, Benjamin EJ. Anger and hostility predict the development of atrial fibrillation in men in the Framingham Offspring Study. *Circulation* 2004 Mar 16;109(10):1267-71.

Eddy DM. Assessment of intensive lifestyle changes for the treatment of coronary heart disease. July 12, 2000. No further details available (reference provided by CMS).

Ellingsen I, Hjermann I, Abdelnoor M, Hjerkin EM, Tonstad S. Dietary and antismoking advice and ischemic heart disease mortality in men with normal or high fasting triacylglycerol concentrations: a 23-y follow-up study. *American Journal of Clinical Nutrition* 2003 Nov; 78(5):935-40.

Elliott-Eller M, Weidner G, Pischke C, Marlin R, Li M, Mendell N, et al. Clinical events in heart disease patients with low left ventricular ejection fraction: 3 year follow-up results. *Proceedings of the International Academy of Cardiology, 3rd World Congress on Heart Disease – New trends in Research, Diagnosis and Treatment*; 2003 July; Washington DC, USA.

Engblom E, Korpilahti K, Hamalainen H, Ronnema T, Puukka P. Quality of life and return to work 5 years after coronary artery bypass surgery. Long-term results of cardiac rehabilitation. *J Cardiopulm Rehabil* 1997;17:29-36.

Esposito K, Giugliano D, Nappo F, Marfella R; Campanian Postprandial Hyperglycemia Study Group. Regression of carotid atherosclerosis by control of postprandial hyperglycemia in type 2 diabetes mellitus. *Circulation* 2004 Jul 13;110(2):214-9. Epub 2004 Jun 14.

Esselstyn CB Jr. Updating a 12-year experience with arrest and reversal therapy for coronary heart disease (an overdue requiem for palliative cardiology). *Am J Cardiol* 1999 Aug 1;84(3):339-41, A8.

Family Heart Study Group. Randomised controlled trial evaluating cardiovascular screening and intervention in general practice: principal results of British family heart study. *BMJ* 1994;308:313-20.

Fields JZ, Walton KG, Schneider RH, Nidich S, Pomerantz R, Suchdev P, et al. Effect of a multimodality natural medicine program on carotid atherosclerosis in older subjects: a pilot trial of Maharishi Vedic Medicine. *Am J Cardiol* 2002 Apr 15;89(8):952-8.

Flanagan DE, Cox P, Paine D, Davies J, Armitage M. Secondary prevention of coronary heart disease in primary care: a healthy heart initiative. *QJM* 1999;92:245-50.

Frasure-Smith N, Prince R. The ischemic heart disease life stress monitoring program: impact on mortality. *Psychosomatic Med* 1985;47:431-45.

Frasure-Smith N, Lesperance F, Gravel G, Masson A, Juneau M, Talajic M, et al. Social support, depression, and mortality during the first year after myocardial infarction. *Circulation* 2000 Apr 25;101(16):1919-24.

Friedman M, Thoreson C, Gill J, et al. Alteration of type A behavior and reduction in cardiac recurrences in post-myocardial infarction patients. *Am Heart J* 1984;108:237-48.

Galatius S, Gustafsson F, Kistorp CM, Nielsen PH, Atar D, Hildebrandt PR. Establishment of a nurse-led heart failure clinic. Design and baseline data from the first two years. *Ugeskr Laeger* 2003;165:686-90.

Geil PB, Anderson JW, Gustafson NJ. Women and men with hypercholesterolemia respond similarly to an American Heart Association step 1 diet. *J Am Diet Assoc* 1995 Apr;95(4):436-41.

George BJ, Goldberg N. The benefits of exercise in geriatric women. *Am J Geriatr Cardiol* 2001 Sep-Oct;10(5):260-3.

Ghoncheh S, Smith JC. Progressive muscle relaxation, yoga stretching, and ABC relaxation theory. *J Clin Psychol* 2004 Jan;60(1):131-6.

Gielen S, Schuler G, Hambrecht R. Exercise training in coronary artery disease and coronary vasomotion. *Circulation* 2001 Jan 2;103(1):E1-6.

Gleason JA, Bourdet KL, Koehn K, Holay SY, Schaefer EJ. Cardiovascular risk reduction and dietary compliance with a home-delivered diet and lifestyle modification program. *J Am Diet Assoc* 2002 Oct;102(10):1445-51.

Gould KL, Ornish D, Kirkeeide R, Brown S, Stuart Y, Buchi M, et al. Improved stenosis geometry by quantitative coronary arteriography after vigorous risk factor modification. *Am J Cardiol* 1992 Apr 1;69(9):845-53.

Gould KL, Ornish D, Scherwitz L, Brown S, Edens RP, Hess MJ, et al. Changes in myocardial perfusion abnormalities by positron emission tomography after long-term, intense risk factor modification. *JAMA* 1995 Sep 20;274(11):894-901.

Grimm RH Jr., for the MRFIT. The Multiple Risk Factor Intervention Trial in the US. A summary of results at four years in special intervention and usual care men. *Preventive Medicine* 1983;12:185-90.

Grundy SM, Cleeman JI, Merz CN, Brewer HB Jr, Clark LT, Hunninghake DB, et al. National Heart, Lung, and Blood Institute; American College of Cardiology Foundation; American Heart Association. Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines. *Circulation* 2004 Jul 13;110(2):227-39.

Gulati M, Pandey DK, Arnsdorf MF, Lauderdale DS, Thisted RA, Wicklund RH, et al. Exercise capacity and the risk of death in women: the St James Women Take Heart Project. *Circulation* 2003 Sep 30;108(13):1554-9.

Hakim AA, Curb JD, Petrovitch H, Rodriguez BL, Yano K, Ross GW, et al. Effects of walking on coronary heart disease in elderly men: the Honolulu Heart Program. *Circulation* 1999 Jul 6;100(1):9-13.

Hambrech R, Walther C, Mobius-Winkler S, Gielen S, Linke A, Conradi K, et al. Percutaneous coronary angioplasty compared with exercise training in patients with stable coronary artery disease: a randomized trial. *Circulation* 2004 Mar 23;109(11):1371-8. Epub 2004 Mar 08.

Harris DE, Record NB, Gipson GW, Pearson TA. Lipid lowering in a multidisciplinary clinic compared with primary physician management. *Am J Cardiol* 1998;81:929-33.

Hedback B, Perk J. 5-year results of a comprehensive rehabilitation programme after myocardial infarction. *Eur Heart J* 1987;8:234-42.

Imperial Cancer Research Fund OXCHECK Study Group. Effectiveness of health checks conducted by nurses in primary care: final results of the OXCHECK Study. *BMJ* 1995;310:1099-1104.

Jain SC, Uppal A, Bhatnagar SO, Talukdar B. A study of response pattern of non-insulin dependent diabetics to yoga therapy. *Diabetes Res Clin Pract* 1993 Jan;19(1):69-74.

Jukema JW, Bruschke AV, van Boven AJ, Reiber JH, Bal ET, Zwinderman AH, et al. Effects of lipid lowering by pravastatin on progression and regression of coronary artery disease in symptomatic men with normal to moderately elevated serum cholesterol levels. The Regression Growth Evaluation Statin Study (REGRESS). *Circulation* 1995 May 15;91(10):2528-40.

Kawachi I, Sparrow D, Vokonas PS, Weiss ST. Symptoms of anxiety and risk of coronary heart disease. The Normative Aging Study. *Circulation* 1994 Nov;90(5):2225-9.

Ketola E, Makela M, Klockars M. Individualised Multifactorial Lifestyle Intervention Trial for High-Risk Cardiovascular Patients in Primary Care. *British Journal of General Practice* 2001 Apr;51(465):291-294.

Koertge J, Weidner G, Billings JH, Elliott-Eller M, Lipsenthal L, Merritt-Worden T, et al. Medical and psychosocial characteristics of CHD patients in the

Multicenter Lifestyle Demonstration Project: Results from a 1-year follow-up [Abstract]. *Psychosomatic Medicine* 2002;64:85-174.

Koertge J, Weidner G, Elliott-Eller M, Scherwitz L, Merritt-Worden TA, Marlin R, et al. Improvement in medical risk factors and quality of life in women and men with coronary artery disease in the Multicenter Lifestyle Demonstration Project. *Am J Cardiol* 2003 Jun 1;91(11):1316-22.

Kornitzer M, De Backer G, Dramaix M, Thilly C. The Belgian Heart Disease Prevention Project. Modification of the coronary risk profile in an industrial population. *Circulation* 1980;61:18-25.

Krachler M, Lindschinger M, Eber B, Watzinger N, Wallner S. Trace elements in coronary heart disease: Impact of intensified lifestyle modification. *Biol Trace Elem Res* 1997 Dec;60(3):175-85.

Kris-Etherton PM, Harris WS, Appel LJ; American Heart Association. Nutrition Committee. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circulation* 2002 Nov 19;106(21):2747-57.

Lampert R, Joska T, Burg MM, Batsford WP, McPherson CA, Jain D. Emotional and physical precipitants of ventricular arrhythmia. *Circulation* 2002 Oct 1;106(14):1800-5.

Lear SA, Ignaszewski A, Linden W, Brozic A, Kiess M, Spinelli JJ, et al. A randomized controlled trial of an extensive lifestyle management intervention (ELMI) following cardiac rehabilitation: study design and baseline data. *Curr Control Trials Cardiovasc Med* 2002 Nov 12;3(1):9.

Lesperance F, Frasure-Smith N, Talajic M, Bourassa MG. Five-year risk of cardiac mortality in relation to initial severity and one-year changes in depression symptoms after myocardial infarction. *Circulation* 2002 Mar 5;105(9):1049-53.

Lewin RJ, Furze G, Robinson J, Griffith K, Wiseman S, Pye M, et al. A randomised controlled trial of a self-management plan for patients with newly diagnosed angina. *Br J Gen Pract* 2002 Mar;52(476):194-6, 199-201.

Lewis CE, Resnik BA. Nurse clinics and progressive ambulatory patient care. *N Engl J Med* 1967;277:1236-41.

Liao XX, Ma H, Dong YG, Tang AL, Tao J, Yang XQ. Effects of early rehabilitative intervention on heart rate variability and quality of life in patients with uncomplicated acute myocardial infarction. *Zhonghua Linchuang Kangfu Zazhi* 2003;7(21):2934-2935.

Lichtenstein AH, Van Horn L. Very low fat diets. *Circulation* 1998 Sep 1;98(9):935-9.

Lindholm LH, Ekblom T, Dash C, Eriksson M, Tibblin G, Schersten B on behalf of the CELL Study Group. The impact of health care advice given in primary care on cardiovascular risk. *BMJ* 1995;310:1105-9.

Maggioni AP. Secondary prevention: improving outcomes following myocardial infarction. *Heart* 2000 Sep;84 Suppl 1:i5-7:discussion i50.

Malach M, Imperato PJ. Depression and acute myocardial infarction. *Prev Cardiol* 2004 Spring;7(2):83-90.

Marra S, Paolillo V, Spadaccini F, Angelino PF. Long-term follow-up after a controlled randomized post-myocardial infarction rehabilitation programme: effects on morbidity and mortality. *Eur Heart J* 1985;6:656-63.

Marshall J, Penckofer S, Llewellyn J. Structured postoperative teaching and knowledge and compliance of patients who had coronary artery bypass surgery. *Heart Lung* 1986;15:76-82.

Matthews KA, Gump BB, Harris KF, Haney TL, Barefoot JC. Hostile behaviors predict cardiovascular mortality among men enrolled in the Multiple Risk Factor Intervention Trial. *Circulation* 2004 Jan 6;109(1):66-70.

Meer H. Coronary heart disease health improvement programme. Final report. Barking and Havering Health Authority, Romford (United Kingdom). Directorate of Public Health, Oct 1999. 67 p. Available from British Library Document Supply Centre-DSC:6966.200(120) In: Public health research report No. 120.

Meland E, Laerum E, Ulvik RJ. Effectiveness of two preventive interventions for coronary heart disease in primary care. *Scand J Prim Health Care* 1997;15:57-64.

Merritt TA, Scherwitz LW, Brown SE, Billings JH, Armstrong WT, Sparler SL, et al. Changes in lifestyle and exercise capacity. *Journal of Cardiopulmonary Rehabilitation* 1990;10(9):354.

Merritt T, Ornish D, Scherwitz L, Billings J, Elliott L, Lipsenthal L, et al. Effects of intensive lifestyle changes on coronary heart disease risk factors and clinical status in self-selected heart patients. *Journal of Cardiopulmonary Rehabilitation* 1995;15(5):353.

Merritt T, Ornish D, Scherwitz L, Billings J, Elliott L, Lipsenthal L, et al. Preliminary results from the multicenter lifestyle trial: Can self-selected heart

patients maintain intensive lifestyle changes? *Journal of Cardiopulmonary Rehabilitation* 1995;15(5):354.

Merritt-Worden TA, Pettengill EB, Ornish D. Gender comparisons in an intensive comprehensive risk factor reduction program [Abstract]. *Journal of Cardiopulmonary Rehabilitation* 2003;23(5):376.

Miettinen TA, Pyorala K, Olsson AG, Musliner TA, Cook TJ, Faergeman O, et al. Cholesterol-lowering therapy in women and elderly patients with myocardial infarction or angina pectoris: findings from the Scandinavian Simvastatin Survival Study (4S). *Circulation* 1997 Dec 16;96(12):4211-8.

Miller ER 3rd, Erlinger TP, Young DR, Jehn M, Charleston J, Rhodes D, et al. Results of the Diet, Exercise, and Weight Loss Intervention Trial (DEW-IT). *Hypertension* 2002 Nov;40(5):612-8.

Mittleman MA, Maclure M, Sherwood JB, Mulry RP, Tofler GH, Jacobs SC, et al. Triggering of acute myocardial infarction onset by episodes of anger. Determinants of Myocardial Infarction Onset Study Investigators. *Circulation* 1995 Oct 1;92(7):1720-5.

National Cholesterol Education Program (NCEP). Detection, evaluation, and treatment of high blood cholesterol in adults (adult treatment panel III). Final Report. Third Report of the National Cholesterol Education Program (NCEP), National Heart, Lung, and Blood Institute, National Institutes of Health, NIH Publication No. 02-5215, September 2002.

Ness AR, Hughes J, Elwood PC, Whitley E, Smith GD, Burr ML. The long-term effect of dietary advice in men with coronary disease: follow-up of the Diet and Reinfarction trial (DART). *European Journal of Clinical Nutrition* 2002 Jun;56(6):512-8.

Nicholson AS, Sklar M, Barnard ND, Gore S, Sullivan R, Browning S. Toward improved management of NIDDM: A randomized, controlled, pilot intervention using a lowfat, vegetarian diet. *Prev Med* 1999 Aug;29(2):87-91.

Oldenburg B, Martin A, Greenwood J, Bernstein L, Allan R. A controlled trial of a behavioral and educational intervention following coronary artery bypass surgery. *J Cardiopulmonary Rehabil* 1995;15:39-46.

Ornish D, Scherwitz LW, Doody RS, Kesten D, McLanahan SM, Brown SE, et al. Effects of stress management training and dietary changes in treating ischemic heart disease. *JAMA* 1983 Jan 7;249(1):54-9.

Ornish D, Brown SE, Scherwitz LW, Billings JH, Armstrong WT, Ports TA, et al. Can lifestyle changes reverse coronary heart disease? The Lifestyle Heart Trial. *Lancet* 1990 Jul 21;336(8708):129-33.

Ornish D. Avoiding revascularization with lifestyle changes: The Multicenter Lifestyle Demonstration Project. *Am J Cardiol* 1998 Nov 26;82(10B):72T-76T.

Ornish D, Scherwitz LW, Billings JH, Brown SE, Gould KL, Merritt TA, et al. Intensive lifestyle changes for reversal of coronary heart disease. *JAMA* 1998 Dec 16;280(23):2001-7.

Ornish D. Statins and the soul of medicine. *American Journal of Cardiology* 2002;89(1):1286-90.

Ornish DM, Pettengill EB. Marked improvements in biomedical and psychosocial cardiac risk factors from a community-based lifestyle change program. *Proceedings of the Annual Meeting of the American Heart Association*, 2003.

Ornish D. Concise Review: intensive lifestyle changes in the management of coronary heart disease. Concise review related to chapter 241: the pathogenesis of atherosclerosis. Printed from Harrison's Online (www.harrisononline.com). McGraw-Hill Companies, 2001-2004 (http://www.accessmedicine.com/server-java/arknoid/amed/harrisons/ex_editorials/ed1209... accessed 09 Dec 2004).

Ornish D. Chapter 8: Intensive life-style changes in the management of coronary heart disease. No further details available (reference provided by CMS).

Ornish D, Hart JA. Chapter 34: Multiple risk factor intervention trials. No further details available (reference provided by CMS).

Pater C, Ditlef Jacobsen C, Rollag A, Sandvik L, Erikssen J, Karin Kogstad E. Design of a randomized controlled trial of comprehensive rehabilitation in patients with myocardial infarction, stabilized acute coronary syndrome, percutaneous transluminal coronary angioplasty or coronary artery bypass grafting: Akershus Comprehensive Cardiac Rehabilitation Trial (the CORE Study). *Curr Control Trials Cardiovasc Med* 2000;1(3):177-183.

Peiss B, Kurlito B, Rubenfire M. Physicians and nurses can be effective educators in coronary risk reduction. *J Gen Intern Med* 1995;10:77-81.

Pettengill EB, Pearson J, Pifalo B, Wilhelm A, Lipsenthal L, Merritt T et al. Community-based lifestyle change program produces substantial reductions in biomedical and psychosocial cardiac risk factors. 4th Scientific Forum on Quality of Care and Outcomes Research in Cardiovascular Disease and Stroke; October 13-14, 2002; Washington, DC.

Pfisterer M, Buser P, Osswald S, Allemann U, Amann W, Angehrn W, et al; Trial of Invasive versus Medical therapy in Elderly patients (TIME) Investigators. Outcome of elderly patients with chronic symptomatic coronary artery disease with an invasive vs optimized medical treatment strategy: one-year results of the randomized TIME trial. *JAMA* 2003 Mar 5;289(9):1117-23.

Picard MH, Schwartz DC, Ahn DK, Kraemer HC, Berger WE, Blumberg R, et al. Cost-benefit analysis of early return to work after uncomplicated acute myocardial infarction. *Am J Cardiol* 1989;63:1308-14.

Pischke C, Weidner G, Billings J, Elliott-Eller M, Lipsenthal L, Merritt-Worden T, et al. Lifestyle changes among heart disease patients with diabetes: results from the multicenter lifestyle demonstration project. *Proceedings of the 62nd Annual Meeting & Scientific Sessions of the American Diabetes Association*; 2002 June 14-18; San Francisco, USA.

Pitt B, Waters D, Brown WV, van Boven AJ, Schwartz L, Title LM, et al. Aggressive lipid-lowering therapy compared with angioplasty in stable coronary artery disease. Atorvastatin versus Revascularization Treatment Investigators. *N Engl J Med* 1999 Jul 8;341(2):70-6.

Pollock ML, Franklin BA, Balady GJ, Chaitman BL, Fleg JL, Fletcher B, et al. AHA Science Advisory. Resistance exercise in individuals with and without cardiovascular disease: benefits, rationale, safety, and prescription: An advisory from the Committee on Exercise, Rehabilitation, and Prevention, Council on Clinical Cardiology, American Heart Association; Position paper endorsed by the American College of Sports Medicine. *Circulation* 2000 Feb 22;101(7):828-33.

Pozen MW, Stechmiller JA, Harris W, Smith S, Fried DD, Voigt GC. A nurse rehabilitator's impact on patients with myocardial infarction. *Medical Care* 1977;15:830-7.

Prochaska JO, Johnson S, Lee P. The transtheoretical model of behavior change. In Shumaker SA, Schron EB, Ockene JK, McBee WL (Eds.), *The handbook of health behavior change – second edition* (pp. 59-84) Springer Publishing Company.

Pyke SDM, Wood DA, Kinmonth AL, Thompson SG, for the Family Heart Study Group. Change in coronary risk and coronary risk factor levels in couples following lifestyle intervention. *Arch Fam Med* 1997;6:354-60.

Rahe RH, Ward HW, Hayes V. Brief group therapy in myocardial infarction rehabilitation: three- to four-year follow-up of a controlled trial. *Psychosomatic Med* 1979;41:229-

Rihal CS, Raco DL, Gersh BJ, Yusuf S. Indications for coronary artery bypass surgery and percutaneous coronary intervention in chronic stable angina: review of the evidence and methodological considerations. *Circulation* 2003 Nov 18;108(20):2439-45.

Roderick P, Ruddock V, Hunt P, Miller G. A randomized trial to evaluate the effectiveness of dietary advice by practice nurses in lowering diet-related coronary heart disease risk. *Br J Gen Pract* 1997;47:7-11.

Roman O, Gutierrez M, Luksic I, Chavez E, Camuzzi AL, Villalon E, et al. Cardiac rehabilitation after acute myocardial infarction. 9-year controlled follow-up study. *Cardiology* 1983;70:223-31.

Rose G, Heller RF, Pedoe HT, Christie DGS. Heart disease prevention project: a randomised controlled trial in industry. *BMJ* 1980;i:747-751.

Rubenstein LV, Kahn KL, Reinisch EJ, Sherwood MJ, Rogers WH, Kamberg C, et al. Changes in quality of care for five diseases measured by implicit review, 1981 to 1986. *JAMA* 1990 Oct 17;264(15):1974-9.

Ruo B, Rumsfeld JS, Hlatky MA, Liu H, Browner WS, Whooley MA. Depressive symptoms and health-related quality of life: the Heart and Soul Study. *JAMA* 2003 Jul 9;290(2):215-21.

[No authors listed] Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet* 1994 Nov 19;344(8934):1383-9.

Schechtman G, Wolff N, Byrd JC, Hiatt JG, Hartz A. Physician extenders for cost-effective management of hypercholesterolemia. *J Gen Intern Med* 1996;11:277-86.

Schneider RH, Staggers F, Alexander CN, Sheppard W, Rainforth M, Kondwani K, et al. A randomised controlled trial of stress reduction for hypertension in older African Americans. *Hypertension* 1995 Nov;26(5):820-7.

Sdringola S, Nakagawa K, Nakagawa Y, Yusuf SW, Boccalandro F, Mullani N, et al. Combined intense lifestyle and pharmacologic lipid treatment further reduce coronary events and myocardial perfusion abnormalities compared with usual-care cholesterol-lowering drugs in coronary artery disease. *J Am Coll Cardiol* 2003 Jan 15;41(2):263-72.

Shaffer J, Wexler LF. Reducing low-density lipoprotein cholesterol levels in an ambulatory care system. Results of a multidisciplinary collaborative practice lipid clinic compared with traditional physician-based care. *Arch Intern Med* 1995;155:2330-5.

Shintani TT, Beckham S, Brown AC, O'Connor HK. The Hawaii Diet: ad libitum high carbohydrate, low fat multi-cultural diet for the reduction of chronic disease risk factors: obesity, hypertension, hypercholesterolemia, and hyperglycemia. *Hawaii Med J* 2001 Mar;60(3):69-73.

Simpson DR, Dixon BG, Bolli, P. Effectiveness of Multidisciplinary Patient Counselling in Reducing Cardiovascular Disease Risk Factors Through Nonpharmacological Intervention: Results From the Healthy Heart Program. *Canadian Journal of Cardiology* 2004 Feb;20(2):177-186.

Sivarajan ES, Newton KM, Almes MJ, Kempf TM, Mansfield LW, Bruce RA. Limited effects of outpatient teaching and counseling after myocardial infarction: a controlled study. *Heart Lung* 1983;12:65-73.

The South-East London Screening Study Group. A controlled trial of multiphasic screening in middle-age: results of the South-East London Screening Study. *Int J Epidemiol* 1977;6:357-63.

Specchia G, De Servi S, Scire A, et al. Interaction between exercise training and ejection fraction in predicting prognosis after a first myocardial infarction. *Circulation* 1996;94:978-82.

Stahle A, Mattsson E, Ryden L, Unden A, Nordlander, R. Improved physical fitness and quality of life following training of elderly patients after acute coronary events. A 1 year follow-up randomized controlled study.[see comment]. *European Heart Journal* 1999 Oct;20(20):1475-84.

Starkey C, Michaelis J, Lusignan S. Computerised systematic secondary prevention in ischemic heart disease: a study in one practice. *Public Health* 2000;114:169-75.

Stern MJ, Cleary P. The national exercise and heart disease project. Long-term psychosocial outcome. *Arch Intern Med* 1982;142:1093-7.

Strandberg TE, Pitkala K, Berglind S, Nieminen MS, Tilvis RS. Multifactorial cardiovascular disease prevention in patients aged 75 years and older: A randomized controlled trial: Drugs and Evidence Based Medicine in the Elderly (DEBATE) Study. *Am Heart J* 2001 Dec;142(6):945-51.

Taddei S, Galetta F, Virdis A, Ghiadoni L, Salvetti G, Franzoni F, et al. Physical activity prevents age-related impairment in nitric oxide availability in elderly athletes. *Circulation* 2000 Jun 27;101(25):2896-901.

Thoresen CE, Friedman M, Gill JK, Ulmer DK. The recurrent coronary prevention project. Some preliminary findings. *Acta Med Scand* 1982;660(Suppl):172-92.

Townsend J, Piper M, Frank AO, Dyer S, North WRS, Meade TW. Reduction in hospital readmission stay of elderly patients by a community based hospital discharge scheme: a randomized controlled trial. *BMJ* 1988;297:544-7.

Tu JV, Pashos CL, Naylor CD, Chen E, Normand SL, Newhouse JP, et al. Use of cardiac procedures and outcomes in elderly patients with myocardial infarction in the United States and Canada. *N Engl J Med* 1997 May 22;336(21):1500-5.

Vale MJ, Jelinek MV, Best JD, Dart AM, Grigg LE, Hare DL, et al; COACH Study Group. Coaching patients On Achieving Cardiovascular Health (COACH): a multicenter randomized trial in patients with coronary heart disease. *Arch Intern Med* 2003 Dec 8-22;163(22):2775-83.

Van Drenth BB, Hulscher EJJ, Mokkink HGA, van de Lisdonk EH, van der Wouden JC, Grol RPTM. Effects of outreach visits by trained nurses on cardiovascular risk-factor recording in general practice. A controlled trial. *Eur J Gen Pract* 1997;3:90-5.

Vedin A, Wilhelmsson C, Tibblin G, Wilhelmsen L. The postinfarction clinic in Goteborg, Sweden. *Acta Med Scand* 1976;200:453-6.

von Birgelen C, Hartmann M, Mintz GS, Baumgart D, Schmermund A, Erbel R. Relation between progression and regression of atherosclerotic left main coronary artery disease and serum cholesterol levels as assessed with serial long-term (> or =12 months) follow-up intravascular ultrasound. *Circulation* 2003 Dec 2;108(22):2757-62.

Wallner S, Watzinger N, Lindschinger M, Smolle KH, Toplak H, Eber B, et al. Effects of intensified lifestyle modification on the need for further revascularization after coronary angioplasty.[see comment]. *European Journal of Clinical Investigation* 1999 May;29(5):372-9.

Wasson J, Gaudette C, Whaley F, Sauvigne A, Baribeau P, Welch G. Telephone care as a substitute for routine clinic follow-up. *JAMA* 1992;267:1788-93.

Waters D, Higginson L, Gladstone P, Kimball B, Le May M, Boccuzzi SJ, et al. Effects of monotherapy with an HMG-CoA reductase inhibitor on the progression of coronary atherosclerosis as assessed by serial quantitative arteriography. The Canadian Coronary Atherosclerosis Intervention Trial. *Circulation* 1994 Mar;89(3):959-68.

Weber F, Barnard RJ, Roy D. Effects of a high-complex-carbohydrate, low-fat diet and daily exercise on individuals 70 years of age and older. *J Gerontol* 1983 Mar;38(2):155-61.

Weidner G, Pischke C, Eller M, Ornish D. Clinical profile and outcome of heart disease patients with low LVEF: results from the Multicenter Lifestyle Demonstration Project. Presented at the annual meeting of the American Psychosomatic Society, Phoenix, AZ, March 5-8, 2003.

Weinberger M, Smith DM, Katz BP, Moore PS. The cost-effectiveness of intensive post-discharge care: a randomized trial. *Med Care* 1988;26:1092-1102.

Weingarten S, Reidinger M, Conner L, Johnson B, Ellrodt AG. Reducing lengths of stay in the coronary care unit with a practice guideline for patients with congestive heart failure: Insights from a controlled clinical trial. *Med Care* 1994;32:1232-43.

Weintraub WS, Clements SD Jr, Crisco LV, Guyton RA, Craver JM, Jones EL, et al. Twenty-year survival after coronary artery surgery: an institutional perspective from Emory University. *Circulation* 2003 Mar 11;107(9):1271-7.

Williams JE, Paton CC, Siegler IC, Eigenbrodt ML, Nieto FJ, Tyroler HA. Anger proneness predicts coronary heart disease risk: prospective analysis from the atherosclerosis risk in communities (ARIC) study. *Circulation* 2000 May 2;101(17):2034-9.

Woollard J, Burke V, Beilin LJ, Verheijden M, Bulsara MK. Effects of a general practice-based intervention on diet, body mass index and blood lipids in patients at cardiovascular risk. *Journal of Cardiovascular Risk* 2003 Feb;10(1):31-40.

Woollard J, Burke V, Beilin LJ. Effects of General Practice-Based Nurse-Counselling on Ambulatory Blood Pressure and Antihypertensive Drug Prescription in Patients at Increased Risk of Cardiovascular Disease. *Journal of Human Hypertension* 2003 Oct; 17(10):689-695.

Yu-Poth S, Zhao G, Etherton T, Naglak M, Jonnalagadda S, Kris-Etherton PM. Effects of the National Cholesterol Education Program's Step I and Step II dietary intervention programs on cardiovascular disease risk factors: a meta-analysis. *Am J Clin Nutr* 1999 Apr;69(4):632-46.